

AD-A095 907

NAVAL DENTAL RESEARCH INST GREAT LAKES IL
A COMPREHENSIVE REVIEW OF NEW ATTACHMENT THERAPY. (U)
JAN 81 M R WIRTHLIN

F/6 6/5


UNCLASSIFIED

NDRI-PR-81-01

NL

1 of 1

004
001907





LEVEL

20


NDRI-PR 81-01
January 1981

1981

AD A 095907

A COMPREHENSIVE REVIEW OF NEW ATTACHMENT THERAPY

M. R. WIRTHLIN



NAVAL
DENTAL RESEARCH
INSTITUTE

Naval Medical Research and Development Command
Bethesda, Maryland

DBG FILE COPY

81 2 2 082

NAVAL DENTAL RESEARCH INSTITUTE
NAVAL BASE, BLDG. 1-H
GREAT LAKES, ILLINOIS 60088

A Comprehensive Review of
New Attachment Therapy

M. R. Wirthlin

Research Progress Report NDRI-PR 81-01
Work Unit 63706N M0095PN003.3010
Naval Medical Research and Development Command
National Naval Medical Center
Bethesda, Maryland 20014

The opinions expressed herein are those of the author and cannot be construed as reflecting the views of the Navy Department or the Naval Service at large. The use of commercially available products does not imply endorsement of the products or preference to other similar products on the market.

This document has been approved for public release; distribution is unlimited.

Approved and released by:



M. R. WIRTHLIN, JR.
Captain, DC, USN
Commanding Officer

Table of Contents

Introduction.....	1
The Tooth Side.....	2
Subgingival Plaque.....	2
Antimicrobial Chemicals.....	2
Calculus.....	2
Zone of Enamel.....	3
Zone of Pathologically Exposed Root Surface.....	3
The Cementum matrix.....	3
Changes in the matrix.....	4
Changes in the mineral.....	4
Changes in the physical state.....	5
Histologic changes.....	6
Adsorption of Toxins.....	6
Detoxification.....	7
Rationale for root planing.....	7
Demineralization.....	8
Zone of the Epithelial Attachment.....	9
Probing the epithelial attachment.....	9
The dental cuticle.....	10
Forming a new junctional epithelium.....	10
The cementum surface.....	11
Zone of the Connective Tissue Attachment.....	11
Repair of the connective tissue attachment.....	12
Cementogenesis.....	12
The Periodontal Side.....	14
Zone of the Marginal Epithelium.....	14
Subgingival curettage--Histologic Studies.....	14
Subgingival curettage--Clinical studies.....	15
Chemicals in curettage.....	15
Excising the pocket epithelium.....	16
Zone of the Gingival Connective Tissue.....	17
Gingival flaps--Histologic studies.....	17
Gingival flaps--Clinical studies.....	18
Gingival grafts.....	19
The periosteum.....	19
Coronal positioning.....	20
Adaptation.....	21
Cell adherence.....	21
Zone of the Bony Defect.....	22
Decortication.....	22
Transseptal fibers.....	22
Curettage--Clinical studies.....	22
Curettage--Histologic studies.....	23
Retarding epithelization.....	24
Implants of Oral Bone--Clinical Studies.....	24
Implants of Oral Bone--Histologic Studies.....	25
Implants of Autogenous Red Marrow and Bone--Clinical Studies.....	25
Implants of Autogenous Red Marrow and Bone--Histologic Studies.....	26
Implants of Allogeneic Marrow and Bone--Clinical Studies.....	27
Implants of Allogeneic Marrow and Bone--Histologic Studies.....	27

Table of Contents - 2

Implants from another species.....	28
Implants of non-bony tissues.....	28
Implants of alloplastic materials.....	29
The Interradicular Zone.....	29
Implants of autogenous oral bone.....	29
Implants of autogenous marrow and bone.....	30
Demineralization.....	30
The Key to Success.....	31
References.....	33
Figures.....	75

New attachment therapy is that which regenerates a more coronal attachment of the periodontal tissues to tooth, after the natural attachment has been destroyed by trauma or disease-- most often the chronic inflammatory periodontal disease, periodontitis. This is distinguished herein from reattachment, or healing by scar,¹ subsequent to surgical detachment. The terms new attachment and reattachment have been used interchangeably by some, and with specific intent by others. "True reattachment" of new cementum, periodontal ligament, and alveolar bone has been distinguished from "soft tissue reattachment" of epithelium and connective tissue coronal to the bony crest,² for instance.

In many cases no goal of new attachment or reattachment has been stated prospectively, at all; and various surgical procedures have developed empirically, based on subjective observations from clinical practice and conceptual speculations about pocket elimination.³ There is a lack of a generally accepted philosophy and no agreement on a goal.⁴ Rather than debate the merits of various "named" surgical procedures it appears time to examine the potential of periodontal tissues, considering the biological processes which bring about healing, and the application of knowledge to clinical benefit.⁵

Following his review of the world literature on periodontal therapy,⁶ Ratcliff¹ proposed an analysis of repair systems in periodontal therapy. Ratcliff¹ stated, "There are four basic forms of periodontal therapy used for elimination of pockets. These are: (1) shrinkage, (2) excision, (3) healing by scar, and (4) new attachment." These were, essentially, accepting the existing dentogingival junction, or cutting it away and growing a new one. Shrinkage was considered the elimination of the edema developed as a part of the inflammatory reaction to irritants, and with a constant position of the epithelial attachment, shrinkage after hygienic procedures would reduce the pocket depth. Excision, such as gingivectomy, and healing by scar, such as with apically repositioned flaps, produce little to no regeneration of a new attachment. For purposes of analysis of new attachment, the tooth side of the periodontal lesion (Figure 1) was considered to have four zones. After the dissection and removal of the pocket lining (Figure 2), the contributions of the zones to new attachment were discussed.¹ Acknowledging Ratcliff's contribution¹ to our understanding, this review will consider other's points of view and later research findings in different zones on the tooth side and the periodontal tissues side, and their applications to new attachment therapy.

THE TOOTH SIDE

SUBGINGIVAL PLAQUE

There is little doubt that the removal of bacterial plaque is essential for the successful healing of a new dentogingival junction. The level of attachment is generally parallel to and congruent with the apical extent of bacterial plaque on the root surface.⁷ The distances from the advancing "front" of the plaque to the nearest periodontal fibers on extracted teeth were measured and found to be 0.3 to 1.5 mm near the cemento-enamel junction but only 0.2 to 1.0 mm nearer the apices.^{8,9} In many cases after new attachment procedures, plaque is inadvertently left on the roots of teeth despite attempts at meticulous instrumentation with overlapping strokes. The removal of plaque is especially difficult if it is hidden in resorption bays in the cementum,^{9,10} at the depth of the pocket, and in the furca.⁹ After meticulous instrumentation of the roots and flushing with sterile saline, bacteria are dispersed on the tooth and soft tissue sides of the pocket, and it may be necessary to use a chemical disinfectant.¹¹

Antimicrobial Chemicals

Pockets have been instilled, after scaling, with emetin,¹² "hydrogen ammonium fluoride," iodine,¹³ "pyrozone" and "hydronaphthol",¹⁴ acriflavin, metaphen, weak iodine or hypertonic salt solution,¹⁵ Dakin's solution,¹⁶ Dunlop oxygen¹⁷ and zinc chloride solution dispersed with electrodes,¹⁸ but without experimental evidence of effectiveness. These treatments were considered to be passing "fads" and not as important as basic scaling, curettage, and oral hygiene.¹⁹ More recently, penicillin has been advocated, because infection was considered the enemy of reinsertion of connective tissue fibers in new cementum.²⁰ The users of strong salt solutions^{21,22} and local delivery of tetracycline by hollow fiber devices^{23,24} have claimed a marked reduction of spirochetes and clinical improvement. There is no support in the literature that local in tasche antimicrobial agents will effect, or assist in, healing by new attachment.

CALCULUS

There is unanimous agreement that calculus must be removed.²⁵⁻²⁷ Despite the report that junctional epithelium may present a normal attachment to disinfected calculus,²⁸ sterilized calculus retains irritants.²⁹ Inadequate removal of calculus will reactivate periodontal disease even with good

supragingival plaque control.^{7,9,10} The removal of calculus is difficult due to its location, hardness, and modes of attachment which may be by cuticles, surface irregularities, penetration of bacteria, undercuts in resorption bays,³⁰⁻³² or penetration between separations of cementum.³³ Root planing enhances the chances of the complete removal of calculus,³⁴ but smooth planed roots may still have deposits.³⁵⁻³⁷ The bacterial flora in a pocket are reduced after subgingival calculus removal;^{24,38,39} and good practice would seem to dictate the removal of calculus, even if a surgical approach was required for access.^{4,40} Past attempts to use acid in pockets to dissolve calculus⁴¹ were thought adverse because experience showed that it made healing slow.⁴²

ZONE OF ENAMEL

Since Ratcliff's analysis,¹ research has shown that epithelium not only readapts to enamel after detachment,⁴³ but that it reforms an attachment which seems identical to that before treatment.^{44,45} The requirements for epithelial attachment to enamel are most likely the absence of accumulated bacteria and their toxins,⁴³ and regenerating junctional epithelium--not oral sulcular or pocket epithelium.⁴⁶ The early strength of the new dentogingival junction in resistance to rupture forces appears to be related to regeneration of epithelial attachment to enamel.⁴⁷

Connective tissue new attachment to enamel has not been reported, but a cementum is found over cervical enamel⁴⁸ which Listgarten⁴⁹ has termed afibrillar cementum. Epithelial reattachment can occur to afibrillar cementum,⁵⁰ and there is one single instance reported of a short connective tissue attachment to cementum on cervical enamel.⁵¹

Enamel projections protrude into furcation areas of multi-rooted teeth.⁵²⁻⁵⁸ Some investigators have found a relationship of enamel projections to pockets;^{57,58} and it has been recommended that enamel projections be ground away in order to provide opportunity for connective tissue new attachment,⁵⁹ even though histology shows they may be covered with cementum.^{55,60}

ZONE OF PATHOLOGICALLY EXPOSED ROOT SURFACE

The Cementum Matrix

The cementum which can become involved in new attachment therapy may be of the acellular, cellular, or intermediate type, depending on the depth of the loss of attachment and the amount of previous instrumentation. Most often, the

acellular cementum in the cervical area is exposed. Even the dentin could be exposed. The cementum has matrix fibers that parallel the root surface^{61,62} and finer fibrils in irregular arrangement^{62,63} with the crossbanding of collagen. The matrix is mineralized and presents alternating dark and light bands on microradiography. The lines are thicker and more irregular in cellular than in acellular cementum.^{62,64} The mineral crystals parallel the collagen fibers,⁶² and mask the collagen matrix.⁶⁵ Principal fibers of the periodontal ligament penetrate the cementum at right angles to the surface and their cores are unmineralized.⁶² In cellular cementum these Sharpey's fibers are not as densely packed as in acellular cementum.⁶² Channels of cementocyte lacunae and their canaliculi are found in cellular cementum.⁶² Intermediate cementum, found near the dentin surface, starts about one-third down the root and has a "collagen hiatus" thought to be protoplasmic inclusions of cells.^{66,67} Between the Sharpey's fibers on the surface are found cell processes, finely granular material, and a three to five micron wide zone of collagen fibers and fine size mineral crystals, the precementum.^{62,68} Repair of cementum may be by acellular or cellular new cementum.^{48,60,61} Dentin intertubular matrix has an irregular arrangement of collagen fibers which are masked by mineral. The dentinal tubules are surrounded by a mineralized peritubular matrix which has no collagen fibers.⁶⁹

Changes in the Matrix

The amino acid composition of the organic matrix of human cementum is that of collagen⁷⁰⁻⁷² with the $\alpha 1(I)_2\alpha 2$ composition typical of other Type I collagens which normally calcify.⁷² No statistically significant differences in collagen composition occur in cementum which is pathologically exposed, except that if affected by dental caries there is loss of the hydroxyproline, proline and glycine typically found in collagen and a gain of amino acids thought to be of bacterial origin.⁷³ The bacterial cell-wall material, alpha, epsilon-diaminopimelic acid is found in carious cementum.⁷³ The amino acid composition of the outer dentin of periodontally diseased teeth may be altered.⁷⁴

Changes in the Mineral

The mineral content of normal cementum changes only slightly from cervical to midroot or apical areas,^{75,76} but the surface of exposed cementum is hypermineralized as revealed by microradiography.^{32,77-83} This hypermineralized zone is usually about 10 microns wide, but is sometimes up to 50 microns deep.^{78,80}

Electron micrographs show crystals on exposed surfaces that are disarranged, larger, or denser.^{81,82,84,85} The denser surface was found to stop at the beginning of the epithelial attachment.⁸¹ When specimens are demineralized, the surface zone often does not present collagen crossbanding in the cementum matrix.^{81-83,85} Analyses of diseased root surfaces show significant increases of calcium,^{78,83,86} magnesium,^{78,86} phosphorus,^{78,83,86} fluoride,^{83,86,87} copper and zinc,⁸⁸ and decreased citrate.⁸⁷

The research indicates a remineralization of the surface of cementum by plaque, once the junctional epithelium separates from the tooth. Scaled exposed root surfaces, whether of cementum or dentin, begin to show microradiopacity by four weeks and it is frequently found at eight weeks.⁸¹ Both the serum in the inflammatory exudate and the saliva could be sources of this mineralization.⁸⁹ Armitage⁹⁰ suggested that if the fluoride were available for release, it may act as an inhibitor to adjacent cells, but that it is likely that fluoride is bound in the crystals. Ratcliff¹ speculated that the hypermineralization made exposed cementum much like enamel, and so was a deterrent to connective tissue new attachment. It is possible for junctional epithelium to attach to hypermineralized root surface.⁹¹

Changes in the Physical State

Removal of the diseased root surface by root planing to produce a hard, smooth surface and to eliminate the necrotic surface layers of cementum has been universally recommended. The microhardness of cementum determined by diamond scratches has been scored 85 to 90, compared to 145 to 165 for dentin and 1600 to 1750 for enamel surface,⁹² but there has been found no statistical difference in the hardness of exposed and unexposed cementum in periodontally-diseased teeth.⁹³⁻⁹⁵

The thickness of cervical cementum has been reported to be as little as 10 microns⁴⁸ to 16 microns.⁹⁶ Cementum normally increases in thickness with age,^{48,97} but periodontally-diseased teeth do not show the same amount of apposition as normal teeth.^{98,99}

The microtopography of normal cementum is finely serrated as mineral crystals project at the insertions of collagen fibrils.⁶² Numerous small, knobby projections are seen as remnants of Sharpey's fiber insertions on the surfaces of extracted teeth.¹⁰⁰⁻¹⁰⁵ Root surfaces exposed by periodontal disease were found significantly rougher than healthy areas.¹⁰⁶ The roots of diseased teeth have more resorptions than the healthy, but these defects are

primarily in the apical area.¹⁰⁷ At the cemento-enamel junction there may be a gap in the cementum 5 to 10% of the time.⁶⁵

Studies with the diffusion of dyes and radioactive materials indicate that there is a "circulation" of tissue fluids into cementum, but that it may decrease in the cervical cementum in older teeth and in periodontally involved human teeth.¹⁰⁸⁻¹¹⁵ Thus "vitality" remains a question of definition, and the same fluid exchange which might contribute to healing might also allow the penetration of bacterial toxins.

Histologic Changes

The cementum and sometimes the surface dentin in pathologically exposed roots of teeth fixed in formalin, decalcified, and cut on a cryostat, present fine granules and clumps of granules which are refractile and colored varying shades of brown when viewed by transmitted light, are white with incident light, and irreversibly cleared in alcohol.¹¹⁶ They are not seen in ground sections, except after decalcification.¹¹⁷ On electron micrographs they appear as vacuole-like formations, singly or in clusters.¹¹⁸ These granules may be related to the amount of calculus¹¹⁹ or in some fashion to bacteria, for pulpless teeth with open, infected chambers present granules in cementum which was still attached to bone by the periodontal ligament.¹²⁰ Human teeth removed because of unmanageable periodontitis-like bone destruction have none, or very limited distribution, of these granules.¹²¹

Bacteria can penetrate into cementum in root caries lesions¹²² and surfaces exposed to plaque.¹²³ Even without bacterial invasion there is thought to be an inhibitory factor in diseased roots, the quantity of which inhibitor may be related to destructive effects and the resistance to destruction, and thus to susceptibility to periodontal disease.¹²⁴ Some such inhibitory factor could also prevent successful new attachment therapy.

Adsorption of Toxins

For many years it has been known that very toxic materials were to be found in periodontal pockets,¹²⁵ and these may adsorb to diseased roots. Bacterial cell wall muramic acid¹²⁶ and other antigens^{127,128} are adsorbed on diseased root surfaces. Periodontally affected human roots leach out a material toxic to epithelial cells in culture.¹²⁹ A phenol-extracted material from cementum of diseased roots was found toxic for fibroblasts in culture.¹³⁰ Aleo, *et al.*,¹³¹ demonstrated a reduced fibroblast adherence to periodontally

diseased root surfaces. Blocks of human diseased roots inhibit osteogenesis in an animal model,^{132,133} and stimulate inflammation when implanted in human submucosal pouches.¹³⁴ Phenol^{36,130,135,136} and acid¹³⁶ extracts of diseased root surface shavings have demonstrated the presence of bacterial lipopolysaccharide endotoxins.

Detoxification

Phenol extraction can increase the adherence of fibroblasts in vitro to diseased root surfaces.¹³¹ Hand scaling³⁶ and ultrasonic instrumentation¹³⁵ can markedly reduce the amount of extractable endotoxin from diseased tooth surfaces, while root planing can reduce the endotoxin to the level of healthy teeth^{36,135} and restore fibroblast adherence in vitro.¹³¹ Root planed blocks of diseased human roots seem to generate somewhat less inflammation than scaled blocks, when observed as submucosal implants.¹³⁴

The depth of endotoxin absorption in diseased roots and the concentration level compatible with new attachment are not known. It may be possible to extract toxic bacterial materials with buffers of some ionic strength¹³⁷ or low pH-salt buffers^{127,128} which are kinder to tissues than phenol and strong acids. Buffers and salts could be used after scaling, and thereby avoid deep and invasive root planing. Extracted human teeth with diseased roots treated with phosphate-buffered physiologic saline after scaling were used as controls for a series of experiments with biological agents, selected from a number of chemicals capable of dissociating or degrading endotoxins.¹³⁸ Sodium deoxycholate followed by human plasma fraction Cohn IV₁ produced a statistically significant increase in adherent fibroblasts, in vitro.¹³⁸ There remains much work to be done to perfect the biological preparation of diseased roots by chemicals and to test them in wound healing experiments for their ability to detoxify roots and abet new attachment therapy.

Rationale for Root Planing

Numerous opinions have been expressed advocating root planing. These have included the points of view that the root surface is pitted and a habitat for bacteria,¹³⁹ that the root surface is dead¹⁴⁰ or necrotic,¹⁴¹ that the pericementum is necrotic,¹⁴² or that remnant Sharpey's fiber ends are putrescent.¹⁴³ A fresh, clean surface has been recommended,^{144,145} which some state should be only in the cementum^{146,147} and others into the dentin.¹⁴⁸⁻¹⁵⁰ Elimination of the biologically unacceptable surface cementum contaminated with substances that can elicit inflammation seems to be a current consensus.¹⁵¹⁻¹⁵³

Root planing to a smooth surface in order to remove calculus will probably

remove most of the cementum.³⁴ The consensus of reports on the results of instruments suggests that curettes leave the smoothest root surface.^{35,37,154-168} However, what feels like a smooth surface to the touch of a sharp explorer may be a grossly rough surface for the first adherent cells of a new attachment. Root planing will not prevent the later accumulation of plaque or calculus, which is related more to the patient's hygiene performance;^{31,166,169,170} but it has been reported that rough tooth surfaces facilitate the retention of bacteria.^{43,171}

As opposed to the point of view of smooth planed roots is the view that healing with new cementum occurs in nicks in the root¹⁷² and that flat surfaces are inhibitory.¹⁷³ The recommendation for a microirregular surface¹⁷⁴ seems supported by only one abstract, a report of an experimental, intentional scarification of cementum after a surgical detachment.¹⁷⁵

Demineralization

Nearly one hundred years ago "aromatic sulfuric acid" was used in deep pockets with claims of remarkable results of new attachment.¹⁷⁶ Claims of new attachment after use of hydrochloric acid¹⁷⁷ lactic acid,^{178,179} salicylic acid¹⁸⁰ and citric acid (used to "neutralize" antiformin)^{181,182} have been made. Black¹⁸³ dismissed the use of acids as "radical" and pointed out that no proponents had demonstrated a connective tissue new attachment by histologic evidence.

Descriptions of histologic repair of surgically-created defects in animals have led to the selection of citric acid as a method for "conditioning" roots and promoting accelerated reattachment.¹⁸⁴⁻¹⁸⁷ Acid etching of normal human cementum exposed the broad bundles and interlacing fibrils of collagen in the cementum matrix.¹⁸⁸ Citric acid applied to diseased root surfaces revealed no collagen fibers, unless the cementum was first removed by root planing.¹⁸⁹ The acid treatment also enlarges the dentinal tubules,¹⁸⁶⁻¹⁸⁹ into which the healing connective tissue enters.^{186,187} The report of Frank, *et al.*,¹⁹⁰ is used as a justification for artificial demineralization. In that study, electron micrographs depict scattered, fine inorganic crystals on the dentin surface among collagen fibers in an area of connective tissue new attachment, one month following a surgical procedure. What seems overlooked is that those investigators, in addition to suggesting that this was a crystal dissolution, also stated that the findings were compatible with beginning of mineralization,¹⁹⁰ or a precementum.

Decalcified normal roots implanted into subcutaneous tissues of rats did not seem to promote connective tissue attachment,¹⁹¹ and when implanted into submucosa of humans led to resorption or exfoliation.¹⁹² The replantation of endodontically-prepared monkey teeth which had been treated with acid caused exfoliation or accentuated resorption and ankylosis, compared to undemineralized controls.¹⁹³

The citric acid conditioning of natural and artificial furcation defects in dogs has demonstrated some remarkable success in new attachment and closure of the defects.^{194,195} The closure of deep pockets in humans has been reported.¹⁹⁶ Histologic study suggests this is mostly by a new, long attachment of junctional epithelium,¹⁹⁷ although some connective tissue adherence and a slight amount of new cementum at the apical extent of the pocket has been demonstrated.¹⁹⁸ In a comparison trial, citric acid conditioned teeth gained 2.1 mm clinical attachment and controls 1.5 mm, six months after modified Widman surgery.¹⁹⁹ Citric acid conditioning has not improved procedures for coverage of denuded roots in humans,^{200,201} but has shown significant new connective tissue attachment in baboons.²⁰²

ZONE OF THE EPITHELIAL ATTACHMENT

The attachment of epithelium to the surface of the tooth provides a biological seal, which, with the attachment of connective tissue elements to the tooth, comprise the dentogingival junction.²⁰³ The occluso-apical length of the junctional epithelium ranges from 0.1 mm to 2.7 mm in the periodontal pocket,²⁰⁴⁻²⁰⁷ and it is quite variable. On extracted teeth it is seen as a "plaque free zone".²⁰⁷⁻²¹¹ It is important to new attachment therapy to know that the junctional epithelium is continually renewed.²¹² Although it has generally been thought that the strength of the dentogingival junction was due to the contribution of connective tissue, the epithelial attachment has surprising strength.^{47,213,214} At the base of the pocket, bacteria are in contact with the junctional epithelium and "pioneer" organisms are found to have advanced into the interstices of deeper cells.²¹⁵ As a result, the junctional epithelium presents characteristics of epithelium over disturbed connective tissue.²¹⁶

Probing the Epithelial Attachment

Most authors agree that the epithelial attachment should be removed to obtain a connective tissue new attachment.^{1,207,217} Thin steel shims inserted in the gingival crevice appear to penetrate to the depth of the epithelial

attachment.²⁰⁷ However, periodontal probes may wedge to a halt at variable positions related to the health of the tissues.^{207,211,218-229} If pre-surgical "initial" therapy is performed to reduce gingival inflammation and dissection then carried out at the bottom of the probable pocket, the junctional epithelial cells may remain on the tooth side as depicted in Figure 2.

Operative dentistry procedures²³⁰ and simple scaling²³¹ may tear or remove the epithelial attachment unintentionally, while intentional procedures to remove it may not completely eliminate it.^{207,232,233} However, the presence of some patches of junctional epithelium may not prevent connective tissue new attachment more coronally.^{234,235}

The Dental Cuticle

Between the junctional epithelium and the tooth surface there is a dental cuticle.^{49,236-243} The dental cuticle may remain on exposed cementum.^{30,81,237,244} The dental cuticle should be removed in attempts at connective tissue new attachment because new cementum may not deposit on it, according to Box²³⁷ and Waerhaug.²⁰⁷

The exact nature of the dental cuticle is not known. Because it appeared hyaline in histological sections it was once thought to be keratin.²⁴⁵ More recently it has been thought to arise as a gradual thickening and condensation of basement laminar material.²⁴⁶ On the basis of stains it has been said to contain hemoglobin,²⁴⁷ protein,²⁴⁸⁻²⁵⁰ and carbohydrates, probably neutral mucopolysaccharides.^{251,252} For others, there is thought to be carbohydrate which remains unidentified.^{249,253,254} Sodium deoxycholate can remove epithelial cells in vitro, and leaves behind this cuticle and attachment substance.²⁵⁵

Dental cuticle was not seen under a new junctional epithelial attachment at two to three weeks after scaling,⁴⁴ but it has been seen at later intervals after reattachment,²⁵⁶ on enamel and calculus after scaling,²⁸ four months after gingivectomy,⁴⁵ and 35 days to a year after flap operations.^{50,257,258}

Forming A New Junctional Epithelium

The epithelial attachment is renewed by the coronal movement of cells,²⁵⁹ and, after detachment, the artificially deepened crevice closes from the bottom, up, like closing a zipper.²⁶⁰ With little to no inflammation present, the junctional epithelium has a low mitotic activity, but it rises dramatically

with inflammation.^{232,261} If some of the junctional epithelium is left on the tooth after a wound of minimal trauma, it participates only slightly in the epithelial regeneration, which is derived primarily from the oral epithelium at the gingival margin.^{47,262}

The Cementum Surface

Underneath the junctional epithelium, its attachment lamina, and the dental cuticle, the cementum surface appears to be varied.⁶⁸ Selvig⁶⁸ found it had a decreased density of the mineral crystals ranging from 0.2 to 18 microns deep, but generally to a depth of less than four microns in four of nine specimens. In other sections the mineral crystals were densely packed at the surface, while subjacent layers were almost devoid of crystals. The loss of mineral crystals revealed the crossbanding of the collagen, except in some regions where the surface decalcified zone failed to exhibit typical collagen structure.⁶⁸ It is most probable that even a light root planing in this zone which would scrape away the epithelial cells and their cuticle would also remove the thin altered surface. Would the exposed collagen fibers in the cementum which still retained their natural crossbanding unite with new collagen fibers of the healing wound? The first fine fibrils of collagen formed on dentin surfaces were generally oriented parallel to the root surface.^{50,187} Whether any "splicing" occurs has not been demonstrated.

ZONE OF THE CONNECTIVE TISSUE ATTACHMENT

The connective tissue attachment is where collagen fibers normally originate in the cementum and pass outward to insert in the gingiva, periosteum, alveolar bone proper, or across the interdental septum to adjacent teeth as transseptal fibers.²⁶³ The distance from the crest of the alveolar bone to the epithelial attachment varies, but is about one millimeter on the average.^{205,206,264}

In this zone, immediately apical to the epithelium, there is an area where the collagenous fibers are found in varying stages of degeneration in the presence of inflammation.^{68,263,265-267} The width of this destruction ranges about 0.5 to 1 mm.^{68,265,266} The fiber dissolution starts with decreased density of fibril packing and longitudinal splitting, progresses to loss of crossbanding, change of orientation, and then fibers appear to be cut off at the cementum surface; but they may retain their crossbanding under the surface.^{68,265} There is a demineralization of the surface of the cementum to a depth of about 0.2 microns in the area of complete fiber dissolution. In the area are found leukocytes, granular debris, and occasional microorganisms.⁶⁸ The fiber dissolution is thought to be enzymatic, as no

multinucleated cells are seen.^{68,267}

The remainder of this zone is of densely packed collagenous fiber bundles and the finely fibrillar indifferent fibers,^{68,268} cells and their processes, and precementum.⁶⁸ The precementum serves to maintain the attachment in health.²⁶⁹⁻²⁷³ Sometimes part of the epithelial network of the former root sheath may be seen.²⁷⁴ Resorptions of the root surface occur as a result of injury, necrosis, or perhaps inflammation, and are normally repaired by cementum to even out the root surface.²⁷⁵⁻²⁷⁹ Sometimes resorption may go deep into the tooth and present an osseous metaplasia repair.²⁸⁰

Repair of the Connective Tissue Attachment

When an incision is made through the gingival crevice or gingival margin to the crest of the alveolar process, the collagen fibers in the zone of connective tissue attachment are severed. When a flap is raised and the root and bone exposed, the fibers are not readily apparent to the eye. Clinicians are repeatedly warned not to scrape them away.^{1,281-283} These fibers have the potential to reattach to the fibers in the repositioned flap. If the root is undamaged, healing by scar does occur.^{47,51,207,282,284-287} If the root in this zone is injured, even with a surgical detachment in normal gingiva, apical migration of the epithelium occurs.^{50,185,186,207,286,288-294} Thus, careless instrumentation may result in an increased loss of connective tissue attachment.

Routine scaling may injure the connective tissue fibers as well as the epithelial attachment.²³¹ The cementoblasts in the area of the connective tissue attachment seem to be very sensitive to trauma, and they "disappear" for a short while after a wound.^{261,295-299} The new cementoblasts may differentiate by some inductive property of the root surface acting on undifferentiated cells,³⁰⁰ but studies seem to indicate a proliferation from the "native cells" on the cementum side of adjacent periodontal ligament.^{287,296,297,301-303} There may also be a migration laterally from adjacent teeth along the transseptal fibers.³⁰⁴ Obtaining a connective tissue reattachment to the root does not ensure bone regeneration.²⁹⁴ The cells, which repopulate the wound, control the type of healing.³⁰⁵ Periodontal ligament cells can produce new cementum and fibers on the root, but bone marrow cells produce resorption and ankylosis of the root.^{296,304-308}

Cementogenesis

When new cementum forms following surgical detachment, it can form on "old" cementum^{50,289,310} or dentin.^{50,289,309-311} New cementum formed in surgical

detachment wounds often follows shallow surface resorption,^{186,285,289,291,292} and is most often found at the apical extent of the wound.^{50,186,285,289,309,310} The earliest material formed is a metachromatic band, probably of mucopolysaccharides,^{50,312} which is later calcified and the collagen-polysaccharide matrix of cementum formed and calcified.^{50,313} The dearth of collagen fibers in that first material would allow artifactual splits in decalcified sections,⁵⁰ but it is not known if this makes the connective tissue reattachment "weaker", in vivo. The earliest collagen fibers of new cementum matrix parallel the root surface.^{50,187,313}

There has been concern that connective tissue healing was impaired on non-vital teeth.^{309,314} Root canal treatment can induce necrosis, resorption and repair of the cementum.³¹⁵ Following surgical detachment,^{316,317} periapical curettage,³¹⁸ or replantation³¹⁹ normal healing was observed histologically on non-vital teeth; while case reports suggest good treatment results for combined endodontic-periodontic lesions.³²⁰⁻³²³ Case reports claim acid treatment is effective in reattachment on non-vital teeth which do not respond to the usual therapy.³²⁴

THE PERIODONTAL SIDE

THE ZONE OF THE MARGINAL EPITHELIUM

With the objective of connective tissue new attachment therapy, the sulcular or pocket epithelium is removed.¹ Scaling with hand instruments and ultrasonic devices may injure the sulcular epithelium by tearing or even removing it, and with tears sometimes extending into the connective tissue.^{231,325-327} With good hygienic measures this simple debridement has been demonstrated histologically to result in reformation of a normal epithelial attachment,^{325,327,328} which may extend almost to the margin of the gingiva.³²⁸

Subgingival Curettage -- Histologic Studies

In subgingival curettage performed as a surgical procedure with the goal of connective tissue new attachment, one attempts, but does not always remove the pocket epithelium.^{207,232,329-335} Remnants may be left at the gingival margin or near the epithelial attachment. The long epithelial rete extensions into the connective tissue may also be left,²³² but if thorough scaling and hygienic measures are used some time prior to curettage, these rete extensions will likely have disappeared.³²⁸ Complete epithelium removal requires many strokes of the curette.³³⁶⁻³³⁸

Following curettage the basal germinative cells of the stratified squamous epithelium at the edge of the wound elongate and migrate as a sheet under the fibrin of the clot, and present evidence of phagocytosis.³³⁹ Depending on the size of the wound and degree of inflammation, complete regeneration may occur in three to 18 days,^{207,222,237,329,331,334,340-343} usually being complete at five to seven days. Excision of the marginal gingiva has been proposed with the rationale of delaying epithelial downgrowth into the wound, thus allowing more time for connective tissue healing, but no experimental evidence confirms that it is more effective.^{335,344-349}

It is conceivable that connective tissue new attachment could result after a well-executed subgingival curettage. Waerhaug reported new cementum in four of 40 human teeth, but thought it was due to injury apical to the base of the pocket.²⁰⁷ Box presented histology of one tooth which showed connective tissue against planed cementum, and an area of new cementum.^{237,350} Schaffer and Zander³⁵¹ presented histologic material demonstrating new connective tissue and epithelial attachment in four cases, in a series where five of eight subjects had clinical pocket reduction after subgingival curettage. Cross³⁵² presented histologic sections of

three teeth in one human with epithelial and connective tissue new attachment. In a study with careful control of hygiene in two monkeys, a long new junctional epithelium was the predominant method of regeneration after root planing and curettage every three months. Occasional "windows" of resorption and connective tissue repair in the epithelium were seen at nine months.²³⁵

Subgingival Curettage -- Clinical Studies

An occasional case report claims considerable reduction of a pocket depth after treatment by subgingival curettage.³⁵⁴ For six patients, one month after subgingival curettage, an average recession of 0.26 mm and 0.9 mm decreased pocket depth were noted.³⁵⁵ In another practice, three months to two years after treatment of 20 patients, one to 7 mm gain of clinical attachments was recorded.³⁴⁵ Significant improvement in 73 humans was recorded in a 36-month study.³⁵⁶ Only five failures were reported in the periodontal maintenance of 103 patients over a six-year period.²⁰ For 157 patients, curettage resulted in a mean 3.0 mm attachment gain.³⁵⁷ Another study, of four-month's duration, showed that subgingival curettage significantly reduced pocket depth and resulted in a gain of attachment for 40 subjects.³⁵⁸ In an experiment on seven dogs treated with curettage six months after surgically creating pockets, 56.7% of the pockets were decreased in depth, with a mean overall gain of attachment of 0.83 mm, 10 months postoperative.³⁵⁹

Short-term surveillance of 44 patients treated with subgingival curettage by the Ramfjord group showed that 62% of the pockets initially deeper than 4 mm were improved, with a mean of 0.2 mm to 0.3 mm gain of attachment level.³⁶⁰ At eight years, data for 78 patients in their long-term study were analyzed on the basis of initial pocket depth. For shallow (1-3 mm) pockets there was about one mm loss of attachment; but in the moderate (4-6 mm) and deep (7-12 mm) pockets there was a sustained gain of attachment and significant reduction of pocket depth.³⁶¹

Chemicals in Curettage

Adjuncts to subgingival curettage have been used empirically until investigations or some years of experience demonstrated if any value existed in continued use. Calcium or sodium sulfide solutions or sodium sulfide and sodium carbonate pastes were once used to "dissolve" epithelium;³⁶²⁻³⁶⁴ but several investigations showed that this very alkaline material was not a specific epithelial solvent, that pockets treated with it healed by epithelial proliferation to former level, and that there was

also connective tissue injury.^{288,365,366} Sodium sulfide was found no more effective than sodium hydroxide.³⁶⁵ Forty percent potassium hydroxide was once used, "neutralized" by hydrochloric acid.⁴¹ Antiformin, a mixture of sodium hypochlorite, sodium carbonate, and sodium hydroxide, was also extremely alkaline, and was "neutralized" by applying citric acid.^{181,182} Likewise, it was found not to dissolve all epithelium and to damage the connective tissue.^{367,368} The use of highly alkaline agents was like an uncontrollable, chemical gingivectomy.³⁶⁹

Phenol, in strengths from 25% to 100%, has been used for an aid in removing epithelium, cuticle, and "infected granulation tissue."^{234,270-273} Interestingly, both phenol and alkali can extract or degrade bacterial endotoxins; but phenol can cause damage to connective tissue³⁷⁴ which, like strong alkalis, would impair new attachment.

The idea of complete removal of epithelium was questioned by Linghorne and O'Connell.^{375,376} In a number of experiments where they interpreted the histology of healing wounds in dogs, they reported that there was a "creeping reattachment" of root resorption, cementum repair and bone regeneration which proliferated coronally at a slow rate, despite the use of procedures which did not attempt to remove pocket epithelium. These results could not be repeated by independent investigators.³⁷⁷ However, one report, where the flap was reflected, the tooth side of the flap coated with gum copal and collodion varnish, autogenous bone from a nearby tooth socket implanted, and the flap closed, showed that connective tissue had formed a new attachment on the tooth, with pocket epithelium nearby in the flap instead of on the tooth surface.³⁷⁸

Excising the Pocket Epithelium

In variations, clinicians have curetted the epithelium before elevation, a gingival or mucoperiosteal flap, have tried to curette the epithelium from the flap after elevation or have dissected the epithelium from the gingival flap by internal bevel incision. The ability of these variations to remove the pocket epithelium and epithelial attachment have not been evaluated to the same extent as has subgingival curettage. A "self-limiting epithelial scalpel"³⁷⁹ was found to effectively remove epithelium on 14% of proximal tissues and about 40% of labial or lingual surfaces.³⁷⁹ Epithelial rete projections in the proximal pocket soft tissue walls between mesial and distal tooth surfaces often come close to touching each other, and so the epithelium can be eliminated completely only by removal of the papilla.³⁸⁰

The early postoperative histology of one study shows that epithelium removal by incision was almost perfect.³⁴¹ It is thought that epithelial removal by incision is less traumatic than curettage and permits better postoperative adaptation to the tooth.³⁸⁰

After dissection, the new epithelium migrates from the cut edge of the marginal epithelium under the clot, and any residual junctional epithelium does not appear to participate in the regeneration.²⁶² The new epithelial attachment has normal ultrastructural appearance on dentin and cementum.^{197, 18} The epithelial new attachment is reformed in about seven days,^{214, 249, 340, 341} at which time it has formed a strong new attachment.²¹⁴ In some areas a 40 nm to 120 nm thick space which may contain amorphous electron dense material is present at four months between epithelium and the root.²⁵⁸

THE ZONE OF THE GINGIVAL CONNECTIVE TISSUE

The residual gingival flap after incision (Figure 1) and the dissection and removal of the lining of the pocket (Figure 2) would ideally leave a consistent wound surface between the epithelium and the alveolar crest.¹ The idealized condition as carried out would appear as in Figure 3. Note that the gingival fibers and the vascular plexus approximately parallel the tooth surface. The thermal effects of ultrasonic curettage may cauterize this surface facing the tooth,^{335, 381-383} or it may be no different than after incision.³⁸⁴ Thermal or chemical coagulation might impede healing.

Hygienic measures and scaling to resolve the gingival inflammation should reduce the possibility of residual ends of pocket epithelium rete in this zone. Those inadvertently left may not constitute a hazard.³⁸⁵ Treatment prior to surgery provides for a regeneration of the free gingival and circular connective tissue fibers.³⁸⁶ Scaling prior to surgery should also reduce hemorrhage, lessen the likelihood of excess granulation tissue in healing, lessen the chance of infection, promote quicker healing, make the tissue firmer and technically easier to incise, and reduce the time required at operation.³⁸⁷⁻³⁹⁰ However, these views are not supported by direct evidence.

Gingival Flaps -- Histologic Studies

A blood clot of fibrin and red blood cells fills the space between the connective tissue and tooth, and is infiltrated with polymorphonuclear leukocytes to form a

"poly band."^{214,299} Granulation tissue proliferates from the periodontal ligament and the flap.^{214,299} The new fibroblasts may come from activation of local fibrocytes, dormant mesenchymal cells, or perhaps from cells of the blood stream.³⁹¹ They migrate into the wound on fibrin strands, followed by capillary buds.³⁹² Interdentally, granulation tissue may also proliferate from marrow spaces. The fibroblasts have a random orientation of polarity on the collagen fibers of the normal gingival connective tissue.³⁹³ With the very close adaptation of a flap having fibers parallel to the tooth, and successful replacement of a thin clot by new fibrous tissue, it is most likely that the result would appear to be what has been called "collagen adhesion."³⁹⁴ There is no evidence that cells of the gingivae have an osteogenic potential and the origin of cementoblasts is uncertain.³⁰¹ It has even been questioned whether gingival connective tissue is able to form anything more than "contact" to a dentin surface.³⁰⁷ Formation of a "collagen adhesion" may prevent or retard the coronal proliferation of periodontal ligament cells which could secure the connective tissue attachment by new cementum. There is no information on the relative strengths of wounds healed by a new attachment with cementum, collagen adhesion, or a long junctional epithelium. It is reasoned that a connective tissue new attachment would give support to the tooth and that a long epithelial attachment could detach and reform a pocket.³⁹⁵

As was found with subgingival curettage, the predominant mode of healing by gingival flaps is a long junctional epithelium new attachment.^{258,299,340,341,342,396-398} Only slight amounts of connective tissue attachment and cementum occur, usually in grooves cut into teeth at the base of the wound.^{199,258,298}

Gingival Flaps -- Clinical Studies

Case reports attest to the clinical success of periodontal disease management by gingival flap procedures.^{16,344,399-402} Clinical studies have followed the results of gingival surgery over short and long-term periods. On 12 patients a postoperative reduction of pocket depth and gingival fluid flow was reported up to four weeks.⁴⁰³ Significant pocket reduction and a slight loss of attachment four months after surgery was reported in a study of 40 patients.³⁵⁸ Over a one-year period, 15 patients were found to have reduction of pocket depth and over one mm gain of attachment.⁴⁰⁴ Another report of 153 pockets in 95 humans demonstrated an average of 3 mm regeneration in 127 pockets, and only seven pockets showed no improvement.⁴⁰⁵ In nine patients, over 2 mm new attachment was found at 12 months,

with a mean of 1.9 mm at three years and 1.5 mm persisting at five years.⁴⁰⁶⁻⁴⁰⁸ Gingival flaps, even without an attempt to return to original preoperative level at the completion of surgery, were found to have a 0.9 mm gain of attachment at six months.⁴⁰⁹

After eight years of clinical trials, the Ramfjord group has found that shallow pockets (1-3 mm) had about one mm loss of attachment, as did those treated by subgingival curettage; but in the moderate (4-6 mm) and deep (7-12 mm) pockets there was a sustained gain of attachment and a significant reduction of pocket depth.³⁶¹

Gingival Grafts

Gingival grafts constitute a use of the zone of connective tissue. In the coverage of denuded roots, no case report or study has documented a new attachment with a free gingival graft.⁴¹⁰⁻⁴¹³ Pedicle grafts moved laterally to cover denuded roots are claimed to be successful in almost every instance.⁴¹⁴⁻⁴¹⁸ A clinical study of pedicle grafts in 15 humans found a significant gain in root coverage.⁴¹⁹ In another study, a significant 3 mm mean gain in attachment was found at six months postoperative in 14 procedures,⁴²⁰ which was sustained at a three-year followup.⁴²¹ Twenty-eight days after creating a denuded root surface surgically, pedicle grafts were performed on 11 dogs in order to study the histology for up to 180 days postoperative. The average result was a 2 mm attachment of epithelium and a 2.1 mm new attachment of connective tissue.⁴²² Two human teeth treated with mucoperiosteal pedicles were seen to have varying amounts of epithelium and connective tissue new attachment.⁴²³ In another report of two cases, the one treated with a split thickness pedicle healed with long junctional epithelium, while one treated with full thickness pedicle had connective tissue and epithelium new attachment.⁴²⁴ Full thickness pedicle grafts have been claimed successful in cases for management of two-walled bony defects,⁴²⁵ furcation problems,⁴²⁶ and adjacent to an edentulous ridge.⁴²⁷

The Periosteum

The success of connective tissue new attachment with mucoperiosteal pedicle grafts moved laterally, as contrasted with replaced gingival flaps, is thought to be associated with the periosteum and its participation in the formation of a union.⁴²⁸

Small fragments of bone often remain on the periosteum elevated from alveolar process.⁴²⁹ New bone formed on the top and bottom of bone spicules which remained attached to periosteum elevated from alveolar process, but little new bone formed where separation had been by elevation through the cambium layer of the periosteum.⁴³⁰ Combination osteoperiosteal flaps produced more new bone when transposed as extraoral pedicle grafts into muscle than did periosteal flaps.⁴³¹ Osteomucoperiosteal pedicle flaps were not successful in forming new attachment to denuded roots in miniature swine.⁴³² Where these flaps were placed over recipient area bone, a new periosteum formed between recipient and donor bone.⁴³²

With the idea that the cambium layer is thin and dormant in the adult, and may be injured when elevated from bone, histologic examination was made of periosteum traumatized by needles jabbed through the gingiva. In three weeks the inflammatory response was followed by proliferation and osteogenesis on the surface of the bone.⁴³³ Case reports of pedicle grafts with "stimulated" periosteum have demonstrated the technique but have not documented the results.^{434,435}

Free grafts of periosteum wrapped around roots during bone implant procedures have not produced a connective tissue new attachment.^{308,436} A descriptive histologic study of free periosteal grafts packed interdentally reported new cementum and connective tissue attachment in the notches marking the depth of the pockets.⁴³⁷ On the basis of case reports, attachment success for free grafts of osteoperiosteal tissues from the tibia has been claimed.⁴³⁸ There has been little work done with pedicle grafts of periosteum.⁴³⁹

Coronal Positioning

Beginning with a gingival flap, then elevating a mucoperiosteal flap, the clinician has the options of replacing the flap or moving it laterally, apically, or coronally. Coronal positioning could bring periosteum onto the root surface with prospects for new attachment and perhaps even increased bone support. Success has been claimed in case reports of coronally positioned mucoperiosteal flaps where the procedure has been started with a gingivectomy⁴⁴⁰ as well as the internal bevel incision.^{441,442} Successful cases of coronally positioned pedicle grafts for coverage of denuded root surfaces of individual teeth have been reported.⁴⁴³⁻⁴⁴⁵ One report has demonstrated 1.43 mm of new attachment 12 months after surgery.⁴⁴⁶ Citric acid treatment of diseased roots in conjunction with coronally positioned pedicles has not shown an

advantage in two human clinical studies.^{200,201} Coronally placed pedicles were found to have no significantly different results from laterally placed pedicles, except that donor teeth for lateral pedicles had slight recession.⁴⁴⁷

Pedicles of mucoperiosteal flaps, elevated after gingivectomy, with their bases at the cuspids and the free ends at midline have been proposed for new attachment,⁴⁴⁸ but others could not obtain the same results.⁴⁴⁹ Limiting such "bridge" pedicles to single teeth, a case report has claimed new attachment, sustained at 18-month follow-up.⁴⁵⁰

Adaptation

An important requirement of new attachment therapy is close adaptation of the gingival tissues to the tooth at the completion of the surgical procedure.⁴⁵¹ When the gingiva is separated by a great amount of clot or debris the healing is delayed.^{262,299} It is a generally held point of view that intimate contact favors a shorter epithelial attachment.^{290,452,453} Close adaptation is thought to result in early formation of an epithelial seal, protecting the connective tissue from bacterial irritants which could interfere with cementogenesis.^{299,454}

Cell Adherence

An area of study with possibly great implications for new attachment therapy is the application of fibronectin in promoting cell adherence to the root surface. Fibronectin is a cell surface glycoprotein which is found on human gingival fibroblasts and provides adherence to culture vessels.⁴⁵⁵ The attachment of gingival fibroblasts to collagen is via the fibronectin, and the attachment probably accounts for the axial orientation of cells to fibers and perhaps even to orientation of new fibrils into bundles.⁴⁵⁶ The attachment of fibroblasts to diseased root surfaces could theoretically be improved by demineralization, thereby exposing collagen for a collagen-fibronectin-cell attachment.⁴⁵⁷ Experiments with non-diseased root surfaces showed that demineralization could double the number of attached fibroblasts in vitro.⁴⁵⁷ The amount of fibronectin in implanted demineralized collagenous bone matrix was found to increase prior to cell proliferation, differentiation, and calcification.⁴⁵⁸ Fibronectin has an affinity for fibrinogen, too, and this may be involved in the locomotion of cells⁴⁵⁹ which migrate into blood clots. There are many exciting new studies which can apply this biological process to new attachment therapy.

THE ZONE OF THE BONY DEFECT

It is thought by some that only when the pocket is within the alveolar process that a new attachment can occur.⁴⁶⁰ Pockets apical to the alveolar crest are classified according to the number of bony walls.⁴⁶¹ The number of bony walls serves as a rational basis for therapy with the three-wall bony defect thought to have the most favorable chance for new attachment therapy.⁴⁶¹

Decortication

In a study of human cadavers, most bony walls observed were of cancellous bone, but sometimes a more sclerotic wall was noted.⁴⁶¹ Cells of osteogenic potential are found in periosteum, endosteum and marrow.⁴⁶² It is thought that perforation of sclerotic bony walls should be made to open into marrow spaces to establish a good blood clot, ensure its rapid revascularization, and facilitate ingress of osteogenic marrow cells.^{463,464} The major repair tissue is from the bone marrow,⁴⁶⁵ and bone resorption would have to open a sclerotic wall before the marrow cells could enter the wound. The arguments for perforation are rational, but there is no experimental evidence of its value.

Transseptal Fibers

Transseptal fibers follow the bony wall surface in interdental angular bony defects.^{466,467} It has been recommended that these fibers be removed, for if left they may be a barrier to proliferation from the marrow.^{461,466,468} Again, this seems rational, but it is done without the support of direct investigation. The granulation tissue which fills the remainder of the defect could have formed from perivascular cells of the gingiva, periosteum, bone marrow, or periodontal ligament. It contains chronic inflammatory cells and the elements of repair; but it is also riddled with elongated epithelial rete,⁴⁶⁹ therefore it is removed.

The number of bony walls is related to the amount of periodontal ligament that is exposed. A three-wall bony defect might have a V or U-shaped exposure of periodontal ligament.⁴⁵⁴ If the usual maximum proliferation outward of the periodontal ligament is about 2 mm, then a complete new attachment is possible in narrow, or in shallow, bony defects.⁴⁵⁴

Curettage -- Clinical Studies

Case reports have indicated a profound response of pocket reduction and apparent bone "fill" on X-ray after treatment of acute periodontal abscesses.⁴⁷⁰⁻⁴⁷⁸ One

point of view holds that the healing is due to a favorable biological state of the root surface following rapid destruction, as opposed to all the changes which occur in chronic pockets with a longer history.⁴⁷⁶ Another view is that there is a radiolucent bony matrix in these lesions, which can be rapidly remineralized as healing occurs.^{470,473} There is really very little evidence in this matter.

Other case reports have claimed "reattachment" and bone fill of chronic defects treated with curettage, as shown by X-ray.⁴⁷⁹⁻⁴⁸⁷ However, X-rays may not accurately depict bony defects or the form of the final results.^{395,488-491} Photography of re-entry operations has demonstrated bone fill.⁴⁸²

Clinical studies with careful documentation have demonstrated gain of attachment upon probing and bone fill in 3-wall, 2-wall, and combination bony defects treated by curettage.⁴⁹²⁻⁴⁹⁸

Curettage -- Histologic Studies

Histologic evaluation of a bony defect in a monkey treated by flap and curettage, and having a 4 mm gain of attachment on probing and bone fill on X-ray, revealed at one year that bone had filled the defect, but that junctional epithelium had migrated apical to the level of planing on the root surface.⁴⁹⁹ In two other animals at one-year postoperative, very little connective tissue new attachment had formed at the base of the defect, and again junctional epithelium was seen on the root in areas of bone fill.⁵⁰⁰ A complete histologic new attachment was observed in 56.3% of 20 monkey bony defects treated by curettage, but about 10% of all specimens had alternating zones of connective tissue attachment and epithelium on the root surface.⁵⁰¹

One human case treated by curettage, which had a shallow crevice on probing and where the X-ray showed bone fill at six months postoperative, disclosed on histology new bone and cementum and a disorganized periodontal ligament.⁵⁰² New bone and cementum were seen in only seven of 21 human block sections in another practice, however two of 21 with no bone were removed at only 14 days postoperative.⁵⁰³ In another study, no bone fill was found in three human blocks removed one year after curettage, and only one specimen had 0.1 mm new cementum, the junctional epithelium having migrated apical to the alveolar crest in all three.⁵⁰⁴

It seems that following curettage, even with probing, X-ray, and re-entry evidence of bone fill, that there may not be a concomitant connective tissue new

attachment. Histology must be done to confirm that epithelium has not migrated to the depth of the former pocket.

Retarding Epithelization

In experiments to determine what could happen if that "bugaboo", epithelium, were excluded, teeth with bony defects were amputated, the defects curetted, and the roots buried by mobilizing flaps.^{397,505,506} Histology at six to eight months usually showed connective tissue new attachments, although the fibers of the new periodontal ligament were parallel to the roots.

Since the epithelium of free gingival grafts usually completely dies and sloughs, it was conceived that bony defects could be covered with free grafts, delaying the migration of epithelium and allowing more time for connective tissue new attachment.⁵⁰⁷ Complete regeneration as determined clinically was found in 60% of 88 human bony defects using this technique, as compared to 40% treated with conventional flaps.⁵⁰⁷ In a histologic study on 20 monkeys, reepithelization was determined to have been retarded by 10 to 12 days by free grafts, and they generally had shorter epithelial attachments.⁵⁰⁸

Implants of Oral Bone -- Clinical Studies

The healing tissue which fills a bony defect could form a dense, fibrous scar, or new bone. It is thought that the matrix of bone contains some inducing or organizing substance and so implanted bone could by its resorption release the stimulus for the formation of new bone in the recipient site.⁵⁰⁹ A number of cases have been reported of successful fill and attachment after the use of autogenous bone implant material obtained from oral donor sites.^{464,510-522}

Clinical studies have quantitated the bone fill seen on X-rays after autogenous bone implants.^{523,524} An increase of over 3 mm mean fill was reported in 44 patients.⁵²⁵ With conventional flap approach, 40% of bony defects obtained complete regeneration in another study.⁵⁰⁷ In eight carefully documented cases, autogenous transplants from oral sites did not appear to markedly influence the outcome, compared to curettage, in 2 and 3-wall defects.⁴⁹⁴ In a study of 91 humans, the same conclusion was made.⁴⁹³ In 25 cases, involving one-wall and furca defects as well as 2 and 3-wall, about 2.9 mm mean bone fill was found with autogenous oral bone implants.⁵²⁵ In a larger group of subjects, it was found that, regardless of the form of the bony defect, there was a statistically significant increase of fill with implants compared to curettage.⁵²⁷ The 2-wall were better when implanted but not the one-wall in another study.⁵²⁸

Implants of Oral Bone -- Histologic Studies

Histologic study has demonstrated that the implantation of autogenous bone chips from an oral site caused bone to regenerate further from the bony defect walls after surgical detachment, than when no implant was used.⁵⁰⁰ Chips smaller than one mm seem to be most effective.⁵²⁹ In epithelized, surgically-created defects the result of autogenous bone implants was "reattachment" in dogs.²⁷⁶ Another study in dogs found histologic evidence of new bone, ligament and cementum in defects implanted, but long epithelial attachment in those not implanted.⁵³⁰ New cementum was seen in grooves at the bottom of defects implanted with fresh autogenous bone from an oral site in another study in dogs.⁵³¹ In 12 monkeys, new cementum, ligament and bone formed in implanted and non-implanted defects.^{501,508} In another study on monkeys both implanted and non-implanted bony defects eventually filled with bone.⁵³²

New cementum formed on old cementum and on dentin, along with new bone formation in one human case at eight months after implanting autogenous cortical and cancellous bone chips.⁵³³ However, the ligament fibers were parallel to the tooth.⁵³³ Another single case report, at 57 months after implanting autogenous bone chips, showed bone fill, but the ligament had considerable artifactual tearing away from the tooth in root planed areas.⁵³⁴ One block section from a clinical study showed new bone, ligament and cementum on a root planed surface which was not denuded of cementum.⁵²⁵ Another instance showed new cementum on old cementum which had been planed, 28 months postoperatively.⁵³⁵ Histologic study of three human teeth at six to 13 weeks after implanting autogenous oral bone showed new bone, a cellular ligament with some functional organization, and new cementum formed on old cementum and on dentin.⁵³⁶ Autogenous bone from oral sites implanted in six human cases when observed at one year showed new bone and cementum in three root planed specimens, but, without root planing, only one of three specimens had new cementum.⁵⁰⁴ There was junctional epithelium on the root apical to the regenerated bone in two of three specimens not root planed, and one of three which were root planed.⁵⁰⁴ A specimen examined 28 weeks after autogenous bone implant was found to have bone regeneration, but with epithelium on the root nearly to the base of the original defect. New cementum was limited to the base of the wound.⁵³⁷

Implants of Autogenous Red Marrow and Bone -- Clinical Studies

Case reports have been presented of the successful use of autogenous cancellous bone and marrow from extraoral sites with the thought that red marrow was the most

osteogenic implant.⁵³⁸⁻⁵⁴³ However, there have also been reports of fresh marrow causing root resorptions of clinical significance.⁵⁴⁴⁻⁵⁴⁷ When using fresh, refrigerated, and frozen red marrow, a clinical study of 52 humans has reported dramatic bone fill in all classes of bony defects.⁵⁴⁸ For others, comparison studies of autogenous red marrow with autogenous cancellous bone from oral sites have demonstrated no significant clinical difference in bony defects.^{494,526} It is clinically significant, however, that autogenous red marrow has resulted in new bone coronal to the alveolar crest, in "overfilled" implant use.^{2,548}

Implants of Autogenous Red Marrow and Bone -- Histologic Studies

Histologic investigation of autogenous red marrow in bony defects in monkeys showed that fresh and frozen marrow from the hip both produced comparable amounts of new bone. The frozen marrow was much better, though, for new cementum and ligament regeneration free of ankylosis.⁵⁰¹ Red marrow implants have resulted in more bone regeneration than non-implanted sites in dogs, but resorption and ankylosis were more common.³⁰³ Fluoride treatment of the donor tissue and recipient site has not affected cementum deposition with red marrow implants in bony defects in dogs.⁵⁴⁹ When composite implants of oral site cancellous bone chips on the root surface and fresh red marrow in the remainder of 2-wall bony defects of monkeys were compared to implants of fresh marrow only, the composite grafts were found to have less resorption and ankylosis. While they often had new cementum on the roots, there was also epithelium apical to the alveolar crest.⁵⁵⁰

New cementum was seen on one extracted tooth from a human clinical trial of marrow implants in 52 patients.⁵⁴⁶ New bone, cementum, and a functionally-oriented ligament were observed on four human teeth in a study of 13 humans with documented pocket reduction and bone fill.² This new cementum formed upon dentin or old cementum, and upon resorbed and unresorbed root surfaces.² The implanted marrow was replaced by new connective tissue at 14 days, and osteogenesis was seen at 24 days, in a series of 15 autogenous bone implants in humans.⁵⁰³ New cementum formed at the level of bone regeneration in narrow or shallow defects; in the deep and wide defects the new bone was more coronal to the new cementum.⁵⁰³ New cementum formed over old cementum or denuded dentin.⁵⁰³ After nine months the implanted bone had been incorporated into new bone or was replaced.⁵⁰³

Implants of Allogeneic Marrow and Bone -- Clinical Studies

Allogeneic bone, preserved in several ways, has been used with success according to case reports.^{436,510,551-554} A clinical study of 20 patients treated with frozen allogeneic bone showed at five to 26 months a mean fill of over 3 mm in all sites.⁵⁵⁵ This was comparable to results with autogenous bone and marrow, including "crestal" apposition.⁵⁵⁵ In another report of eight patients, about 1.5 mm mean bone fill was seen at re-entry one year postoperative.⁵⁵⁶ Freeze-dried allogeneic bone implants in 350 patients were reported to have complete or greater than 50% fill in 60% of the implant sites re-entered.^{557,558} Another report of nine patients also found greater than 50% fill in 60% of sites implanted with freeze-dried bone, but this was not different than the results in control defects treated by curettage.⁵⁵⁹ Decalcified, freeze-dried allogeneic bone implants were found in three cases to have 4 mm to 10 mm new bone produced in defects re-entered at 32 to 104 weeks.⁵⁶⁰

Implants of Allogeneic Marrow and Bone -- Histologic Studies

Histologic observation of fresh allogeneic red marrow implants in surgical defects in 12 dogs showed new bone and cementum at four and eight weeks after implantation.⁵⁶¹ Resorption and repair at the base of bony defects were seen in four dogs, whether the sites received merthiolate-preserved allogeneic bone chips or not.⁵⁶² In six dogs, new bone and cementum attachment were observed in histologic sections of merthiolate-preserved allogeneic bone implanted into surgically-created bony defects.⁴³⁶ One human case of an implant from a bone bank into an intrabony pocket, but which became infected and was removed, showed new bone forming on and between the implanted chips at 3½ months.⁵⁶³ New bone and connective tissue but no cementum were seen on histology of a 2-month postoperative specimen implanted with a frozen allograft of red marrow.⁵⁶⁴ In a companion report, new cementum was seen on old cementum at the base of the bony defect of a one-month specimen.⁵⁵⁵ Fresh and frozen allogeneic red marrow implants in four monkeys were found both to result in new bone, cementum and functionally-oriented ligament fibers.⁵⁶⁵ New cementum and connective tissue attachment were seen in five of six human frozen allogeneic red marrow implants at six months or one year, but junctional epithelium was also seen on the tooth apical to the regenerated bone in four of the six specimens.⁵⁰⁴ Freeze-dried bone implants were seen on histology to have formed new bone at six weeks postoperative, but the results in four dogs were not different than defects treated by curettage.⁵⁶⁶ Decalcified, allogeneic bone matrix was replaced by new

bone in 27 dogs, but at 13 weeks there was ankylosis of new bone to the roots.⁵⁶⁷ In a human case, a biopsy at 17 weeks showed new bone forming on decalcified freeze-dried bone, but no root was available to determine if there was cementogenesis.⁵⁶⁰

Implants from Another Species

Case reports of xenografts, or implants of non-human bone, have claimed good results in a number of instances.^{510,538,568-572} Clinical studies have shown some pocket reduction, attachment gain and bone fill,^{492,573,574} but in comparison to curettage there was essentially no difference.⁴⁹²

Histologic examination showed that bone powder from sheep implanted in six dogs resulted in new bone, ligament and cementum after surgical detachment.⁵⁷⁵ Boiled cow bone powder in dogs appeared to hasten reattachment in surgically-created defects.⁵⁷⁶ A human block section removed 15 years after the implant of boiled cow bone into an 8 mm bony defect showed new cementum and connective tissue on an old, irregular cementum surface presumed to be the former exposed root.⁵⁷⁷ Commercially-produced cow bone was found to be not completely replaced after six months in surgically created defects in six dogs.⁵⁷⁸ Decalcified, despeciated cow bone was found to be incorporated in host bone faster than undemineralized implants when placed under periosteum on alveolar process of dogs.⁵⁷⁹ Anorganic bone has been found to be quite inert.^{590,581} Defatted, deproteinated ox bone was found to be no different than oral autogenous bone chips in 12 monkeys.⁵⁸²

Implants of Non-Bony Tissues

Implants of collagen have not shown significant differences from controls in regeneration.⁵⁸³⁻⁵⁸⁶ Using dentin and cementum chips from human teeth as implants in surgically-created defects in six dogs, epithelium was found on the tooth apical to the new bone in two specimens.⁵⁸⁷ The implantation of cementum and dentin shavings into surgical defects of animals has resulted in osteoid formation on the smaller chips and new cementum,^{588,589} but at four months some areas of ankylosis had occurred.⁵⁸⁹ Cartilage allografts have been claimed to result in new attachment in human cases,^{590,591} and the histology of cartilage implants showed they were replaced by bone in four monkeys.⁵⁹² However, a comparison in dogs showed no difference in repair whether implanted with cartilage or not.⁵⁶² Implanted human sclera has been reported to reduce pockets in human cases.^{593,594} Histologic study of sclera implants in human bony defects shows they are incorporated into host fibrous connective

tissue but are not replaced by bone,^{595,596} and with no cementogenesis at one year.⁵⁹⁶

Implants of Alloplastic Materials

Plaster of Paris implants into surgically created bony defects in 10 dogs were resorbed by six weeks, but showed no more bone than unimplanted defects.⁵⁹⁷ No bone was found at six months in 10 defects in five humans when plaster was mixed and inserted in bony defects as a paste;⁵⁹⁸ but when implanted as set plaster pellets 79% of the defects in 35 cases demonstrated "fill" on X-rays.⁵⁹⁹ Tricalcium phosphate ceramic materials implanted into surgically-created bony defects in dogs were slowly resorbed by multinucleated giant cells, which may have retarded its replacement with bone, but defects were repaired at 22 to 24 weeks.⁶⁰⁰⁻⁶⁰² In six humans, one ceramic implant was exfoliated at three weeks, but in the other five bone fill was seen on X-ray at two to 16 months.⁶⁰³ Compared to allogeneic bone implants, ceramic was found to be no different.⁵⁵⁶ Zirconium oxide ceramic cloth implanted in surgical defects had bone form on its surface in monkeys and dogs.⁶⁰⁴ In five human bony defects, four pyrolytic graphite implants were exfoliated and there was no regeneration in the other.⁶⁰⁵ No new attachment has been found with these materials.

THE INTERRADICULAR ZONE

Implants of Autogenous Oral Bone

The furcae within multi-rooted teeth have always been a difficult management problem in periodontics. The implantation of autogenous bone from oral donor sites into surgically-created furcal defects in dogs resulted in a variable amount of bone regeneration and new cementum in the root nicks marking the base of the defects, while unimplanted controls had bone loss and epithelium grew down into the furcae.⁵³¹ Oral cancellous bone implants in 17 monkeys developed total closure of the furcae in 44% of sites, with new cementum in 34%. Controls with no implants had only four of 27 sites (15%) with total closure and new cementum.⁶⁰⁶ Devitalized oral cancellous bone resulted in no apparent difference than fresh oral cancellous bone in 19 monkeys.⁶⁰⁷ Oral cancellous bone implanted in six dogs did not result in complete regeneration. Epithelium migrated into the furcae of 33 out of 36 sites, despite the flaps having been sutured up on the crowns of the teeth.⁶⁰⁸ No difference was found between use of oral cancellous bone and defatted, deproteinated ox bone. Few sites developed new cementum on the "roof" of the furca.⁵⁸²

Implants of Autogenous Marrow and Bone

Autogenous red marrow implanted into the furcae of two human molars was found to have resulted in complete fill on re-entry at five to six months.⁵³⁹ Fresh red marrow implants in monkeys formed total closure in 12 of 24 sites (50%) and new cementum in 17%, while frozen red marrow had complete closure in 30% and new cementum in 28%. However, 75% of the fresh marrow sites had resorption and ankylosis.⁶⁰⁶

Demineralization

Significantly more bone fill was found in citric acid-treated furcae in eight dogs, compared to the controls, for 13 of 23 sites had complete connective tissue new attachment with cementum and eight of 23 sites had connective tissue attachment but no new cementum, at six weeks.¹⁹⁴ In another study, 27 of 35 sites had complete cementum regeneration and five partial regeneration in furcae of dogs at 12 weeks.⁶⁰⁹ Comparing coronal flap positioning with closure to the preoperative level in conjunction with citric acid treatment of furcae in dogs showed no difference in a third study.⁶¹⁰ The overall results of the three studies was a complete closure in 71% of sites.⁶¹⁰ A preliminary report has indicated new attachment in natural defects as well as surgical defects in dogs.¹⁹⁵ Comparable investigations in humans have not been reported.

As was found with treatment by curettage, bony defects treated with implants of various materials may have probing, X-ray, and re-entry evidence of bone fill, and still may not have a connective tissue new attachment. Only histologic evidence can confirm that epithelium has not migrated apically to its former depth. It is generally thought that bone implants are not needed in "narrow" defects, but there is no study which has established "narrow" limits. Implants have not really demonstrated an inductive effect, but the results with many cases suggest that some particulate implants may be conductive. That is, they may facilitate the migration of osteogenic tissue into the defect. In a similar fashion, do implants of small bone chips provide space for coronal migration of periodontal ligament cells? What is the limit of the migration, and how predictable is it? These questions have not been answered by investigative evidence. Implants of bone do facilitate the regeneration of host bone in sites with low potential for innate regeneration, such as one-wall bony defects, supracrestal and furcae.^{611,612} However, there is still no conclusive evidence that bone implants increase the predictability of connective tissue new attachment.^{613,614}

THE KEY TO SUCCESS

It is a prevalent point of view, eloquently stated by Zander, et al.,⁶¹⁵ that surgical pocket elimination is an essential part of therapy; that to maintain hygiene and the integrity of the attachment there must be re-established "normal" sulcular dimensions. Pocket elimination surgery does not necessarily improve plaque control,⁶¹⁶ however. Another view is that it really may be the therapist's choice of the procedures to use in new attachment therapy; as long as good oral hygiene and maintenance care are ensured, teeth can be maintained in health and function even with a long junctional epithelium, or collagen adhesion, and without new bone fill.⁶¹⁷⁻⁶²⁰ The question is, can a dentition survive in that status?

Patients who receive routine periodontal maintenance every three months do not exhibit a dramatic or prolonged improvement in the absence of personal oral hygiene.⁶²¹ Personal oral hygiene can rapidly reverse gingival inflammation in persons with gingivitis,⁶²² and significantly alter the flora of plaque.⁶²³ Those persons with periodontal pockets need both frequent professional care and personal oral hygiene. Subgingival scaling can have a dramatic effect on the flora of pockets.^{24,39} Scaling and optimal plaque control can resolve inflammation and reduce pocket depth.⁶²⁴⁻⁶²⁸ Scaling, root planing and oral hygiene can cause significant gains in clinical attachment as determined by periodontal probing.^{627,629-631} Scaling and hygiene can even perform as well on attachment gain as subgingival curettage or other surgical procedures.⁶³²⁻⁶³⁵

Inflammation in the healing wound is a primary factor in determining the resultant clinical attachment. Epithelium will most likely extend to the depth of the inflammation.^{234,530} Patients with poor personal hygiene performance before new attachment therapy were found to not have a significant gain of attachment clinically.⁶³⁶ Those persons with poor hygiene postoperatively were found to have a loss of attachment.^{496,637,638} Even with careful experimental control of hygiene, the junctional epithelium may migrate to the depth of root planing, but the health of the junctional epithelium and connective tissue in the new dentogingival junction will be resistant to probing.^{499,500} With control of hygiene and the prevention of postoperative inflammation, significant bone regeneration occurs.^{495,496,498,639,640} Even in the difficult to manage furcation, postoperative hygiene can result in significant bone regeneration.^{641,642}

Long-term case histories and clinical studies of patients treated for periodontal disease attest to the importance of professional maintenance and personal hygiene

by the patient.^{152,361,408,635,643-649} According to Feinstein,⁶⁵⁰ clinicians constantly perform "experiments", the experimental material being a person, prepared by nature, who has much to say about who the investigator will be, and the time, place, and the goals of the experiment. The clinician observes data such as signs and symptoms of illness, demographic, morphologic, and microbiologic evidence. His purpose is to repeat a success of the past with well established procedures. The enumeration of case successes is as scientific as the titration with a biuret or weighing with an analytical balance. Thus these long-term case histories support the value of periodontal therapy. A deepened sulcus or a long epithelial attachment can be considered an acceptable result as long as there is no observation of data which show disease recurrence.

One bit of evidence that the clinician cannot readily gather is the chemical state of the exposed root surface. The laboratory experiments in this area have not determined all the deleterious effects of root exposure nor the most beneficial methods with which to revitalize that surface. It is hoped that patients will not have to shed too much blood in clinical "experiments" before the answers are provided.⁶

It may be that, before long, chemicals or biologic agents will be used as adjuncts to the steel of curettes and scalpels. An antibacterial agent to complete the "toilet of the cavity" after scraping away plaque and calculus would probably help prevent infection. An agent that would dissociate and bind up endotoxin, or a buffer solution that would elute adsorbed toxins and antigens, appear to be sound, innovative and rational. Application of surface binding agents may improve the "plating efficiency" of connective tissue cell adherence, and perhaps prevent the apical migration of epithelium. In these experimental developments, it seems well to remember, "First, do no harm," and to avoid harsh, caustic chemicals.

The current status of new attachment therapy, which seems to be supported by sound prior research, is that clinicians who employ it accomplish a new dentogingival junction of a long epithelial attachment, backed up by healthy collagenous connective tissue, which is functional and maintainable for a long time. The key to success is the attention to hygienic measures.

REFERENCES

1. Ratcliff, P. A.: An analysis of repair systems in periodontal therapy. Periodont Abstr 14: 57, 1966.
2. Dragoo, M. R., and Sullivan, H. C.: A clinical and histological evaluation of autogenous iliac bone grafts in humans: Part I. Wound healing 2 to 8 months. J Periodontol 44: 599, 1973.
3. Waerhaug, J.: Needs and potentials for clinical research in periodontology. J Periodontol 40: 155, 1969.
4. Waerhaug, J.: Review of Cohen: "Role of periodontal surgery." J Dent Res 50: 219, 1971.
5. Melcher, A. H.: In Melcher, A. H., and Bowen, W. H. (eds), Biology of the Periodontium, p 524. New York, Academic Press, 1969.
6. Ratcliff, P. A.: Periodontal therapy-- Review of the literature. In Ramfjord, S. P., Kerr, D. H., and Ash, M. M. (eds), World Workshop in Periodontics, p 277. Ann Arbor, American Academy of Periodontology and University of Michigan, 1966.
7. Waerhaug, J.: A method for evaluation of periodontal problems on extracted teeth. J Clin Periodontol 2: 160, 1975.
8. Waerhaug, J.: Subgingival plaque and loss of attachment in periodontosis as evaluated on extracted teeth. J Periodontol 48: 125, 1977.
9. Waerhaug, J.: Healing of the dentogingival junction following subgingival plaque control. II. As observed on extracted teeth. J Periodontol 49: 119, 1978.
10. Waerhaug, J.: Healing of the dento-epithelial junction following subgingival plaque control. I. As observed in human biopsy material. J Periodontol 49: 1, 1978.
11. Oshrain, H. I., Salkind, A., and Mandel, I. D.: Bacteriologic studies of periodontal pockets ten minutes after curettage. J Periodontol 43: 685, 1972.
12. Beall, E. N.: The present status of emetin in the treatment of pyorrhea. Dent Cosmos 58: 406, 1916.
13. Smith, T. S.: Inflammatory diseases of the gingival margin and periodontal membrane (pyorrhea alveolaris). Dent Cosmos 58: 1239, 1916.
14. Hart, A. C.: Evaluation of decay No. II. Dent Cosmos 43: 201, 1901.
15. Bunting, R. W.: The control and treatment of pyorrhea by subgingival surgery. JADA 15: 119, 1928.
16. Kirkland, O.: The suppurative periodontal pus pocket; its treatment by the modified flap operation. JADA 18: 1462, 1931.

17. Box, H. K.: Oxygen Insufflation in Periodontal Diseases. Springfield, C. C. Thomas, 1955.
18. Sturridge, E.: The inflammatory tissues of the gingival margin and periodontal membrane: Treatment by ionic medication. Dent Cosmos 58: 403, 1916.
19. Smith, R.: Fads in periodontal treatment that have passed into disuse and fundamentals for successful treatment that remain. JADA 26: 1318, 1939.
20. Carranza, F. A.: A technic for reattachment. J Periodontol 25: 272, 1954.
21. Keyes, P. H., Wright, W. E., and Howard, S. A.: The use of phase-contrast microscopy and chemotherapy in the diagnosis and treatment of periodontal lesions--an initial report (I). Quint Internat p 51, 1978.
22. Keyes, P. H., Wright, W. E., and Howard, S. A.: The use of phase-contrast microscopy and chemotherapy in the diagnosis and treatment of periodontal lesions--an initial report (II). Quint Internat p 69, 1978.
23. Goodson, J. M., Haffajee, A., and Socransky, S. S.: Periodontal therapy by local delivery of tetracycline. J Clin Periodontol 6: 83, 1979.
24. Lindhe, J., Heijl, L., Goodson, J. M., and Socransky, S. S.: Local tetracycline delivery using hollow fiber devices in periodontal therapy. J Clin Periodontol 6: 141, 1979.
25. Ramfjord, S. P., and Ash, M. M. Jr.: Periodontology and Periodontics. p 362. Philadelphia, W. B. Saunders, 1979.
26. Moskow, B. S., Baer, P. N., Hazen, S. P., and Turesky, S. S.: In our opinion. What is the role of subgingival calculus in the etiology and progression of periodontal disease? J Periodontol 41: 283, 1970.
27. Nabers, C. L., Prichard, J. F., Robinson, R. E., and Schallhorn, R. G.: In our opinion. In order to secure "fill" in osseous defects is it necessary or desirable to "plane" the exposed root surfaces until they are hard and smooth to the touch? J Periodontol 41: 419, 1970.
28. Listgarten, M. A., and Ellegaard, B.: Electron microscopic evidence of a cellular attachment between junctional epithelium and dental calculus. J Periodont Res 8: 143, 1973.
29. Allen, D., and Kerr, D.: Tissue response in the guinea pig to sterile and nonsterile calculus. J Periodontol 36: 121, 1965.
30. Zander, H. A.: The attachment of calculus to root surfaces. J Periodontol 24: 16, 1953.
31. Kopczyk, R. A., and Conroy, C. W.: The attachment of calculus to root planed surfaces. Periodontics 6: 78, 1968.

32. Selvig, K. A.: Attachment of plaque and calculus to tooth surfaces. J Periodont Res 5: 8, 1970.
33. Moskow, B. S.: Calculus attachment in cemental separations. J Periodontol 40: 125, 1969.
34. Schaffer, E. M.: Histologic results of root curettage of human teeth. J Periodontol 27: 296, 1956.
35. Jones, S. J., Lozdan, J., and Boyde, A.: Tooth surfaces treated in situ with periodontal instruments. Brit Dent J 132: 57, 1972.
36. Jones, W. A., and O'Leary, T. J.: The effectiveness of in vivo root planing in removing bacterial endotoxin from the roots of periodontally involved teeth. J Periodontol 49: 337, 1978.
37. Walker, S. L., and Ash, M. M.: A study of root planing by scanning electron microscopy. Dent Hygiene 50: 109, 1976.
38. Simonton, F. V.: The most significant findings of the California Stomatological Research Group in the study of pyorrhea. J Dent Res 8: 235, 1928.
39. Listgarten, M. A., Lindhe, J., and Hellden, L.: Effect of tetracycline and/or scaling on human periodontal disease. J Clin Periodontol 5: 246, 1978.
40. Ramfjord, S. P., and Ash, M. M. Jr.: Periodontology and Periodontics, p 314. Philadelphia, W. B. Saunders, 1979.
41. Hecker, F.: Pyorrhea Alveolaris. St. Louis, C. V. Mosby, 1913.
42. Smith, T. S.: The successful scientific treatment of periodontal diseases (pyorrhea alveolaris). Dent Items Interest 38: 437, 1915.
43. Waerhaug, J.: Effect of rough surfaces upon gingival tissue. J Dent Res 35: 323, 1956.
44. Thilander, H., and Hugoson, A.: The border zone tooth-enamel and epithelium after periodontal treatment. Acta Odont Scand 28: 147, 1970.
45. Listgarten, M. A.: Electron microscopic features of the newly formed epithelial attachment after gingival surgery. J Periodont Res 2: 46, 1967.
46. Schroeder, H. E., and Listgarten, M. A.: Fine Structure of the Developing Epithelial Attachment of Human Teeth, p 104. Basel, S. Karger, 1971.

47. Wirthlin, M. R., Yeager, J. E., Hancock, E. B., and Gaugler, R. W.: The healing of gingival wounds in miniature swine. J Periodontol 51: 318, 1980.
48. Kronfeld, R.: The biology of cementum. JADA 25: 1451, 1938.
49. Listgarten, M. A.: Electron microscopic study of the gingivo-dental junction of man. Amer J Anat 119: 147, 1966.
50. Listgarten, M. A.: Electron microscopic study of the junction between surgically denuded root surfaces and regenerated periodontal tissues. J Periodont Res 7: 68, 1972.
51. Kohler, C. A., and Ramfjord, S. P.: Healing of gingival mucoperiosteal flaps. Oral Surg 13: 89, 1960.
52. Masters, D. H., and Hoskins, S. W.: Projection of cervical enamel into molar furcations. J Periodontol 35: 49, 1964.
53. Grewe, J. W., Meskin, L. H., and Miller, T.: Cervical enamel projections: Prevalence, location, and extent; with associated periodontal implications. J Periodontol 36: 460, 1965.
54. Leib, A. M., Berdon, J. K., and Sabes, W. R.: Furcation involvements correlated with enamel projections from the cemento-enamel junction. J Periodontol 38: 330, 1967.
55. Moskow, B. S.: Some observations on radicular enamel. J Periodontol 42: 92, 1971.
56. Tsatsas, B., Mandi, F., and Kerani, S.: Cervical enamel projections in the molar teeth. J Periodontol 44: 312, 1973.
57. Bissada, N. F., and Abdelmalek, R. G.: Incidence of cervical enamel projections and its relationship to furcation involvement in Egyptian skulls. J Periodontol 44: 583, 1973.
58. Larato, D. C.: Some anatomical factors related to furcation involvements. J Periodontol 46: 608, 1975.
59. Ramfjord, S. P., and Ash, M. M. Jr: Periodontology and Periodontics, p 666. Philadelphia, W. B. Saunders, 1979.
60. Kerr, D. A.: The cementum: Its role in periodontal health and disease. J Periodontol 32: 183, 1961.
61. Thoma, K. H., and Goldman, H. M.: The pathology of dental cementum. JADA 26: 1943, 1939.
62. Selvig, K. A.: The fine structure of human cementum. Acta Odont Scand 23: 423, 1965.

63. Romaniuk, K.: Some observations of the fine structure of human cementum. J Dent Res 46: 152, 1967. Austral. Sect., IADR Abstr No. 27.
64. Soni, N. N., Van Huysen, G., and Swenson, H. M.: A microradiographic and X-ray, densitometric study of cementum. J Periodontol 33: 372, 1962.
65. Hopewell-Smith, A.: Concerning human cementum. J Dent Res 2: 59, 1920.
66. Blackwood, H. J. J.: Intermediate cementum. Brit Dent J 102: 345, 1957.
67. El Mostehy, M. R., and Stallard, R. E.: Intermediate cementum. J Periodont Res 3: 24, 1968.
68. Selvig, K. A.: Ultrastructural changes in cementum and adjacent connective tissue in periodontal disease. Acta Odont Scand 24: 459, 1966.
69. Johansen, E.: Microstructure of enamel and dentin. J Dent Res 43: 1007, 1964.
70. Rodriguez, M. S., and Wilderman, M. N.: Amino acid composition of the cementum matrix from human molar teeth. J Periodontol 43: 438, 1972.
71. Eastoe, J. E.: Composition of the organic matrix of cementum. J Dent Res 54(Special Issue A): L137, 1975. IADR Abstr L547.
72. Christner, P. J., and Robinson, P. J.: Characterization of human cementum collagen. J Dent Res 54(Special Issue A): 67, 1975. AADR Abstr No. 89
73. Cloutier, P., and Johansen, E.: Studies on organic matrix of sound, exposed and carious cementum. IADR Prog Abstr: 148, 1971. IADR Abstr No. 383.
74. Emslie, R. D., and Stack, M. V.: A chemical change in cervical dentine in periodontal disease. Dent Pract 8: 19, 1957.
75. Selvig, K. A., and Selvig, S. E.: Mineral content of human and seal cementum. J Dent Res 41: 624, 1962.
76. Neiders, M. E., Eick, J. D., Miller, W. A., and Leitner, J. W.: Electron probe microanalysis of cementum and underlying dentin in young permanent teeth. J Dent Res 51: 122, 1972.
77. Forsberg, A., Lagergren, C., and Lönnerblad, O. D.: Dental calculus. A biophysical study. Oral Surg 13: 1051, 1960.
78. Selvig, K. A., and Zander, H. A.: Chemical analysis and microradiography of cementum and dentin from periodontally diseased human teeth. J Periodontol 33: 303, 1962.
79. Yamamoto, H., Sugahara, N., and Yamada, N.: Histopathological and microradiographic study of the exposed cementum from periodontally diseased human teeth. Tokyo Med Dent Univ Bull 13: 407, 1966.
80. Furseth, R., and Johansen, E.: A microradiographic comparison of sound and carious human dental cementum. Archs Oral Biol 13: 1197, 1968.

81. Selvig, K. A.: Biologic changes at the tooth-saliva interface in periodontal disease. J Dent Res 48: 846, 1969.
82. Furseth, R.: Further observations on the fine structure of orally exposed and carious human dental cementum. Archs Oral Biol 16: 71, 1971.
83. Selvig, K. A., and Hals, E.: Periodontally diseased cementum studied by correlated microradiography, electron probe analysis and electron microscopy. J Periodont Res 12: 419, 1977.
84. Herting, H. C.: Electron microscope studies of the cementum surface structures of periodontally healthy and diseased teeth. J Dent Res 46: 1247, 1967. Eur. Div. IADR Abstr No. 18.
85. Yamada, N.: Fine structure of exposed cementum in periodontal disease. Tokyo Med Dent Univ Bull 15: 409, 1968.
86. Wirthlin, M. R., Pederson, E. D., Hancock, E. B., Lamberts, B. L., and Leonard, E. P.: The hypermineralization of diseased root surfaces. J Periodontol 50: 125, 1979.
87. Stepnick, R. J., Nakata, T. M., and Zipkin, I.: The effects of age and fluoride exposure on fluoride, citrate and carbonate content of human cementum. J Periodontol 46: 45, 1975.
88. Sherman, D. B., Ruben, M. P., Goldman, H. M., Breck, F., and Healey, M.: Laser spectrochemical microanalysis of cementum. IADR Prog Abstr: 59, 1966. IADR Abstr No. 85.
89. Stewart, R. T., and Ratcliff, P. A.: The source of components of subgingival plaque and calculus. Periodont Abstr 14: 102, 1966.
90. Armitage, G. C.: Alterations in exposed human cementum. Periodont Abstr 25: 60, 1977.
91. Listgarten, M. A., and Sullivan, H., unpublished data cited by Listgarten, M.: In Stahl, S. S. (ed), Periodontal Surgery, p 189. Springfield, IL., C. C. Thomas, 1976.
92. Hodge, H. C., and McKay, H.: The microhardness of teeth. JADA 20: 227, 1933.
93. Emslie, R. D., and Stack, M. V.: The microhardness of roots of teeth with periodontal disease. Dent Pract 9: 101, 1958.
94. Rautiola, C. A., and Craig, R. G.: The microhardness of cementum and underlying dentin of normal teeth and teeth exposed to periodontal disease. J Periodontol 32: 113, 1961.

95. Warren, E. B., Hansen, N. M., Swartz, M. L., and Phillips, R. W.: Effects of periodontal disease and of calculus solvents on microhardness of cementum. J Periodontol 35: 505, 1964.
96. Skillen, W. G.: Histological and pathological studies of the periodontal membrane, cementum, and dentin with reference to infections. J Nat Dent Assn 8: 621, 1921.
97. Zander, H. A., and Hürzeler, B.: Continuous cementum apposition. J Dent Res 37: 1035, 1958.
98. Hürzeler, B., and Zander, H. A.: Cementum apposition in periodontally diseased teeth. Helv Odont Acta 3: 1, 1959.
99. Müller, G., and Zander, H. A.: Cementum of periodontally diseased teeth from India. J Dent Res 39: 385, 1960.
100. Benson, L. A.: The surface of cementum. J Periodontol 30: 126, 1959.
101. Heuser, H.: Influence of function on human surface cementum. Dent Abstr 8: 145, 1963.
102. Landay, M., Heidere, J., and Ritman, L.: A scanning electron microscopic study of human cementum. IADR Prog Abstr: 62, 1971. IADR Abstr No. 37.
103. Landay, M. A.: A scanning electron microscopic study of human cementum exposed by periodontal disease. IADR Prog Abstr: 56, 1972. IADR Abstr No. 36.
104. Kerry, G. J.: New method for viewing root surfaces. J Periodontol 46: 673, 1975.
105. Sottosanti, J. S.: A possible relationship between occlusion, root resorption, and the progression of periodontal disease. Periodont Abstr 25: 69, 1977.
106. Stoller, S. M.: The microtopography of teeth. J Periodontol 36: 485, 1965.
107. Harvey, B. L. C., and Zander, H. A.: Root surface resorption of periodontally diseased teeth. Oral Surg 12: 1439, 1959.
108. Cahn, L. R.: A preliminary report on the dentinal-cemental communication, with special reference to the abnormally large channels seen in pyorrhetic teeth. Dent Item Int 48: 477, 1926.
109. Stewart-Ross, W. The permeability of the cementum and its reaction to irritation. Brit Dent J 55: 177, 1933.
110. Stones, H. H.: The permeability of cementum. Brit Dent J 56: 273, 1934.
111. Bodecker, C. F., and Lefkowitz, W.: Concerning the "vitality" of the calcified dental tissues. J Dent Res 16: 463, 1937.

112. Blayney, J. R., Wasserman, F., Groetzinger, G., and DeWitt, T. G.: Further studies in mineral metabolism of human teeth by the use of radioactive phosphorus. J Dent Res 20: 559, 1941.
113. Bartlestone, H. J.: Survey of the use of radioactive isotopes in dentistry. N Y Dent J 20: 322, 1950.
114. Wainwright, W. W.: The permeability of human dental root structures to radioactive iodine. J Periodontol 23: 95, 1952.
115. Linden, L-A.: Microscopic observations of fluid flow through cementum and dentine. Odont Revy 19: 367, 1968.
116. Bass, C. C.: A previously undescribed demonstrable pathologic condition in exposed cementum and the underlying dentine. Oral Surg 4: 641, 1951.
117. Armitage, G. C., and Christie, T. M.: Structural changes in exposed human cementum. I. Light microscopic observations. J Periodont Res 8: 343, 1973.
118. Armitage, G. C., and Christie, T. M.: Structural changes in exposed human cementum. II. Electron microscopic observations. J Periodont Res 8: 356, 1973.
119. Benson, L. A.: A study of pathologic condition in exposed cementum. Oral Surg 16: 1137, 1963.
120. Armitage, G. C.: Structural changes in cementum from non-vital human teeth. J Dent Res 53(Special Issue): 80, 1974. IADR Abstr No. 102.
121. Durso, P. J.: Unpublished material.
122. Furseth, R., and Johnsen, E.: The mineral phase of sound and carious human dental cementum studied by electron microscopy. Acta Odont Scand 28: 305, 1970.
123. Sottosanti, J. S., and Garrett, J. S.: A rationale for root preparation--a scanning electron microscopic study of diseased cementum. J Periodontol 46: 628, 1975. Abstr.
124. Morris, M. L.: A new hypothesis of periodontal homeostasis. J Periodontol 51: 49, 1980.
125. Graham, J. W.: The toxicity of sterile filtrate from parodontal pockets. Proc Roy Soc Med 30: 1165, 1937.
126. Armstrong, W. G.: Amino-acid composition of the acquired pellicle of human tooth enamel. Nature 210: 197, 1966.

127. Everhart, D. L., and Stahl, S. S.: Reaction of component recovered from cementum exposed to periodontal pocket with human serum. J Dent Res 56: 538, 1977.
128. Bravman, R. J., Everhart, D. L., and Stahl, S. S.: A cementum-bound antigen: Its reaction with serum antibody and localization, in situ. J Periodontol 50: 656, 1979.
129. Hatfield, C. G., and Baumhanners, A.: Cytotoxic effects of periodontally involved surfaces of human teeth. Archs Oral Biol 16: 465, 1971.
130. Aleo, J. J., DenRenzis, F. A., Farber, P. A., and Varboncouer, A. P.: The presence and biologic activity of cementum-bound endotoxin. J Periodontol 45: 672, 1974.
131. Aleo, J. J., DenRenzis, F. A., and Farber, P. A.: In vitro attachment of human gingival fibroblasts to root surfaces. J Periodontol 46: 639, 1975.
132. Morris, M. L.: The subcutaneous implantation of periodontally diseased roots. J Periodontol 43: 737, 1972.
133. Morris, M. L.: An inhibitory principle in the matrix of periodontally diseased roots. J Periodontol 46: 33, 1975.
134. Lopez, N. J., Belvederessi, M., and de la Sotta, R.: Inflammatory effects of periodontally diseased cementum studied by autogenous dental root implants in humans. J Periodontol 51: 582, 1980.
135. Nishimine, D., and O'Leary, T. J.: Hand instrumentation versus ultrasonics in the removal of endotoxins from root surfaces. J Periodontol 50: 345, 1979.
136. Fine, D. H., Morris, M. L., Tabak, L., and Cole, J. D.: Preliminary characterization of material eluted from the roots of periodontally diseased teeth. J Periodont Res 15: 10, 1980.
137. Söder, P.-O., and Otteskog, P.: Autotransplantation of periodontally involved teeth with use of tissue culture techniques. J Dent Res 54(Special Issue): L70, 1975. IADR Abstr No. L281.
138. Wirthlin, M. R., and Hancock, E. B.: Biological preparation of diseased root surfaces. J Periodontol 51: 291, 1980.
139. Hartzell, T. B.: The practical surgery of the root surface in pyorrhea. Dent Cosmos 53: 513, 1911.
140. Fish, E. W.: Local and remote sequelae of infection in the parodontal sulcus. Proc Roy Soc Med 30: 1149, 1937.
141. Stillman, P. R.: The management of pyorrhea. Dent Cosmos 59: 405, 1917.

142. Merritt, H. H.: The pathology, etiology, and treatment of pyorrhea. Dent Cosmos 60: 574, 1918.
143. Leonard, H. J.: Periodontia treatment for maximum reattachment. JADA 22: 602, 1935.
144. Chace, R.: Methods and values of tooth planing in periodontal therapy. J Periodontol 32: 233, 1961.
145. Chace, R.: Subgingival curettage in periodontal therapy. J Periodontol 45: 107, 1974.
146. Smith, T. S.: Constructive treatment of diseased paradental tissues. JADA 22: 1477, 1935.
147. Bell, D. G.: Indication and contraindication for subgingival curettage in the treatment of paradentosis. J Periodontol 8: 55, 1937.
148. Morris, M. L.: Reattachment of periodontal tissue. A critical study. Oral Surg 2: 1194, 1949.
149. Riffle, A. B.: The cementum during curettage. J Periodontol 23: 170, 1952.
150. Carranza, F. A. Jr.: Glickman's Clinical Periodontology, 5th ed. p 782. Philadelphia, W. B. Saunders, 1979.
151. Robinson, P. J.: Possible roles of diseased cementum in periodontitis. J Prev Dent 2: 3, May-June, 1975.
152. Garrett, J. S.: Root planing: A perspective. J Periodontol 48: 553, 1977.
153. Ramfjord, S. P., and Ash, M. M. Jr.: Periodontology and Periodontics, p 363. Philadelphia, W. B. Saunders, 1979.
154. Barnes, J. E., and Schaffer, E. M.: Subgingival root planing: A comparison using files, hoes and curettes. J Periodontol 31: 300, 1960.
155. Bjorn, H., and Lindhe, J.: The influence of periodontal instruments on the tooth surface. Odont Revy 13: 355, 1962.
156. Burke, S. W., and Green, E.: Effectiveness of periodontal files. J Periodontol 41: 39, 1970.
157. Clark, S. M., Grupe, H. E., and Mahler, D. M.: The effect of ultrasonic instrumentation on root surfaces. J Periodontol 39: 135, 1968.
158. Ewen, S. J.: A photomicrographic study of root scaling. Periodontics 4: 273, 1966.
159. Ewen, S. J., and Gwinnett, A. J.: A scanning electron microscopic study of teeth following periodontal instrumentation. J Periodontol 48: 92, 1977.

160. Green, E., and Ramfjord, S. P.: Tooth roughness after subgingival root planing. J Periodontol 37: 396, 1966.
161. Green, E.: Root planing with sharp and dull curettes. J Periodontol 39: 348, 1968.
162. Johnson, W. N., and Wilson, J. R.: The application of the ultrasonic dental unit to scaling procedures. J Periodontol 28: 264, 1957.
163. Kerry, G. J.: Roughness of root surfaces after use of ultrasonic instruments and hand curettes. J Periodontol 38: 340, 1967.
164. Lie, T., and Meyer, K.: Calculus removal and loss of tooth substance in response to different periodontal instruments. J Clin Periodontol 4: 250, 1977.
165. Pameijer, C. H., Stallard, R. E., and Hiep, N.: Surface characteristics of teeth following periodontal instrumentation: A scanning electron microscope study. J Periodontol 43: 628, 1972.
166. Rosenberg, R. M., and Ash, M. M.: The effect of root roughness on plaque accumulation and gingival inflammation. J Periodontol 45: 146, 1974.
167. Thebaud, J.: Some microscopic aspects of the curetted surface of the cementum after subgingival curettage. Canad Dent A J 17: 127, 1951.
168. Wilkinson, R. F., and Maybury, J. E.: Scanning electron microscopy of the root surface following instrumentation. J Periodontol 44: 559, 1973.
169. Graham, C. J.: Home care effectiveness upon planed teeth and scaled teeth following surgery. J Periodontol 37: 43, 1966.
170. Volpe, A. R., Manhold, J. H., and Hazen, S. P.: In vivo calculus assessment: Part I. A method and its examiner reproducibility. J Periodontol 36: 292, 1965.
171. Swartz, M. L., and Phillips, R. W.: Comparison of bacterial accumulations on rough and smooth enamel surfaces. J Periodontol 28: 304, 1957.
172. Morris, M. L.: The implantation of human dentin and cementum into the subcutaneous tissues of the rat. Periodontics 5: 113, 1967.
173. Morris, M. L.: The effects of root shape and biology on bone growth. J Periodontol 49: 33, 1978.
174. Ruben, M. P.: Significance of root surface preparation in periodontal therapy. Quint Internat 9: 57, 1978.
175. Glickman, I., and Lazansky, J. P.: Reattachment of the marginal gingiva and periodontal membrane in experimental animals. J Dent Res 29: 659, 1950. IADR Abstr No. 11.

176. Marshall, J. S.: A remarkable case of pyorrhea alveolaris, with reproduction of bone, occurring in the practice of Dr. Allport, Chicago, Ill. JAMA 1: 641, 1883.
177. Stewart, H. T.: Partial removal of cementum and decalcification of tooth in the treatment of pyorrhea alveolaris. Dent Cosmos 41: 617, 1899.
178. Younger, W. J.: Pyorrhea alveolaris, from a bacteriological standpoint, with a report of some investigations, and remarks on the treatment. Internat Dent J 20: 413, 1899.
179. Good, R.: Pyorrhea. Dent Digest 12: 355, 1906.
180. Barkmann, L.: A new painless treatment in periodontal disease. Dent Cosmos 73: 674, 1931.
181. Box, K. F.: Concerning the role of antiformin-citric acid in pocket therapy. Ont Dent A J 29: 437, 1952.
182. Hugill, R. A.: A general practitioner's observations on the antiformin-citric acid therapy. Ont Dent A J 29: 441, 1952.
183. Black, A. D.: Treatment of periodontal inflammation coordinated with physiological reactions of the tissues. JADA 24: 1587, 1937.
184. Register, A. A.: Bone and cementum induction by dentin, demineralized in situ. J Periodontol 44: 49, 1973.
185. Register, A. A., and Burdick, F. A.: Accelerated reattachment with cementogenesis to dentin, demineralized in situ. I. Optimum range. J Periodontol 46: 646, 1975.
186. Register, A. A., and Burdick, F. A.: Accelerated reattachment with cementogenesis to dentin, demineralized in situ. II. Defect repair. J Periodontol 47: 497, 1976.
187. Ririe, C. M., Crigger, M., and Selvig, K. A.: Healing of periodontal connective tissues following surgical wounding and application of citric acid in dogs. J Periodont Res 15: 314, 1980.
188. Tonge, C. H., and Boulton, E. H.: Collagen in human cementum as shown by electron microscopy. Nature 182: 459, 1958.
189. Garrett, J. S., Crigger, M., and Egelberg, J.: Effects of citric acid on diseased root surfaces. J Periodont Res 13: 155, 1978.
190. Frank, R., Fiore-Donno, G., Cimasani, G., and Matter, J.: Ultrastructural study of epithelial and connective gingival reattachment in man. J Periodontol 45: 626, 1974.

191. Morris, M. L.: The effect of root decalcification on the formation of "functionally" oriented collagen fibers. J Periodontol 51: 171, 1980.
192. Morris, M. L.: The submucosal implantation of human allogeneic decalcified dentin. J Periodontol 49: 36, 1978.
193. Nordenram, A., Bang, G., and Anneroth, G.: A histopathologic study of replanted teeth with superficially demineralized root surfaces in Java monkeys. Scand J Dent Res 81: 294, 1973.
194. Crigger, M., Bogle, G., Nilvéus, R., Egelberg, J., and Selvig, K. A.: The effect of topical citric acid application on the healing of experimental furcation defects in dogs. J Periodont Res 13: 538, 1978.
195. Bogle, G., Adams, D., Crigger, M., Klinge, B., and Egelberg, J.: New attachment after surgical treatment and acid conditioning of roots in naturally occurring periodontal disease in dogs. J Clin Periodontol 7: 344, 1980. Abstr.
196. Register, A. A.: Human pocket reattachment to dentin, demineralized in situ. J Dent Res 54(Special Issue): 64, 1975. IADR Abstr No. 80.
197. Stahl, S. S., and Froum, S. J.: Human clinical and histologic repair responses following the use of citric acid in periodontal therapy. J Periodontol 48: 261, 1977.
198. Cole, R. T., Crigger, M., Bogle, G., Egelberg, J., and Selvig, K. A.: Connective tissue regeneration to periodontally diseased teeth. J Periodont Res 15: 1, 1980.
199. Cole, R., Nilvéus, R., Ainamo, J., Bogle, G., Crigger, M., and Egelberg, J.: Effect of citric acid application on healing after modified Widman surgery. J Clin Periodontol 7: 345, 1980. Abstr.
200. Otomo, J. A., and Sims, T. N.: Effects of citric acid demineralization on coronally repositioned flaps. J Dent Res 58(Special Issue A): 347, 1979. IADR Abstr No. 1021.
201. Liu, W. J., and Solt, C. W.: A surgical procedure for the treatment of localized gingival recession in conjunction with root surface citric acid conditioning. J Periodontol 51: 505, 1980.
202. Yukna, R. A.: Evaluation of citric acid root preparation during laterally positioned flap procedures. J Dent Res 59(Special Issue A): 451, 1980. AADR Abstr No. 729.
203. Marfino, N. R., Orban, B. J., and Wentz, F. M.: Repair of the dento-gingival junction following surgical intervention. J Periodontol 30: 180, 1959.

204. Ritchey, B., and Orban, B.: The periodontal pocket. J Periodontol 23: 199, 1952.
205. Stanley, H. R.: The cyclic phenomenon of periodontitis. Oral Surg 8: 598, 1955.
206. Wade, A. B.: Instruments. I. Biologic considerations. Dent Pract 7: 193, 1957.
207. Waerhaug, J.: The gingival pocket: Anatomy, pathology, deepening and elimination. Odont Tidskr 60: Suppl 1, 1952.
208. Hoffman, I. D., and Gold, W.: Distances between plaque and remnants of attached periodontal tissues on extracted teeth. J Periodontol 42: 29, 1971.
209. Brady, J. M.: A plaque free zone on human teeth--scanning and transmission electron microscopy. J Periodontol 44: 416, 1973.
210. Saglie, R., Johansen, J. R., and Tollefsen, T.: Plaque-free zones on human teeth in periodontitis. J Clin Periodontol 2: 190, 1975.
211. Powell, B., and Garnick, J. J.: The use of extracted teeth to evaluate clinical measurements of periodontal disease. J Periodontol 49: 621, 1978.
212. Löe, H.: Physiological aspects of the gingival pocket. Acta Odont Scand 19: 387, 1961.
213. Toller, J. R.: The organic continuity of the dentine, the enamel, and the epithelial attachment in dogs. Brit Dent J 67: 443, 1939.
214. Hiatt, W. H., Stallard, R. E., Butler, E. D., and Badgett, B.: Repair following mucoperiosteal flap surgery with full gingival retention. J Periodontol 39: 11, 1968.
215. Saglie, R.: A scanning electron microscopic study of the relationship between the most apically located subgingival plaque and the epithelial attachment. J Periodontol 48: 105, 1977.
216. Ten Cate, A. R.: The dento-gingival junction. An interpretation of the literature. J Periodontol 46: 475, 1975.
217. Carranza, F. A. Jr: Glickman's Clinical Periodontology, 5th ed. p 846. Philadelphia, W. B. Saunders, 1979.
218. Sivertson, J. F., and Burgett, F. J.: Probing of pockets related to the attachment level. J Periodontol 47: 281, 1976.
219. Listgarten, M. A., Mao, R., and Robinson, P. J.: Periodontal probing and the relationship of the probe tip to periodontal tissues. J Periodontol 47: 511, 1976.

220. Spray, J. R., Garnick, J. J., Doles, L. R., and Klawitter, J. J.: Microscopic demonstration of the position of periodontal probes. J Periodontol 49: 148, 1978.
221. Ezis, I., and Burgett, F.: Probing related to attachment levels on recently erupted teeth. J Dent Res 57(Special Issue A): 307, 1978. IADR Abstr No. 932.
222. Armitage, G. C., Svanberg, G. K., and Löe, H.: Microscopic evaluation of clinical measurements of connective tissue levels. J Clin Periodontol 4: 173, 1977.
223. Robinson, P. J., and Vitek, R. M.: The relationship between inflammation and resistance to probe penetration. J Periodont Res 14: 239, 1979.
224. van der Velden, U.: Influence of periodontal health on probing depth and bleeding tendency. J Clin Periodontol 7: 129, 1980.
225. Garnick, J. J., Spray, J. R., Vernino, D. M., and Klawitter, J. J.: Demonstration of probes in human periodontal pockets. J Periodontol 51: 563, 1980.
226. Jansen, J., Pilot, T., and Corba, N.: Histologic evaluation of probing in experimentally induced periodontal defects in beagle dogs. J Clin Periodontol 7: 343, 1980. Abstr.
227. Listgarten, M. A.: Periodontal probing: What does it mean? J Clin Periodontol 7: 165, 1980.
228. Lawther, G. C.: Relationships of periodontal instruments to soft tissues of periodontal pockets. Thesis, University of Minnesota, Minneapolis, 1957.
229. Hancock, E. B., and Wirthlin, M. R.: The location of the periodontal probe tip in health and disease. J Periodontol (in press).
230. Bodecker, C. F., and Lefkowitz, W.: Gingival reattachment to the enamel and its relation to operative dentistry. Dent Cosmos 77: 1106, 1935.
231. Ramfjord, S., and Kiester, G.: The gingival sulcus and the periodontal pocket immediately following scaling of teeth. J Periodontol 25: 167, 1954.
232. Stone, S., Ramfjord, S. P., and Waldron, J.: Scaling and gingival curettage. A radioautographic study. J Periodontol 37: 415, 1966.
233. Weinreb, M. M.: The epithelial attachment. J Periodontol 31: 186, 1960.
234. Ramfjord, S. P.: Experimental reattachment in rhesus monkeys. J Periodontol 22: 67, 1951.

235. Caton, J. G., and Zander, H. A.: The attachment between tooth and gingival tissues after periodic root planing and soft tissue curettage. J Periodontol 50: 462, 1979.
236. Bodecker, C. F., and Applebaum, E.: The clinical importance of the gingival crevice. Dent Cosmos 76: 1127, 1934.
237. Box, H. K.: Treatment of the Periodontal Pocket. Toronto, University Toronto Press, 1928.
238. Glickman, I., and Bibby, B. G.: The existence of cuticular structures on human teeth. J Dent Res 22: 91, 1943.
239. Bass, C. C.: A demonstrable line on extracted teeth indicating the location of the outer border of the epithelial attachment. J Dent Res 25: 401, 1946.
240. Schroeder, H. F.: Ultrastructure of the junctional epithelium of the human gingiva. Helv Odont Acta 13: 65, 1969.
241. Schroeder, H. E., and Listgarten, M. A.: Fine Structure of the Developing Epithelial Attachment of Human Teeth, p 56. Basel, S. Karger, 1971.
242. Listgarten, M. A.: Structure of surface coatings on teeth. A review. J Periodontol 47: 139, 1976.
243. Kobayashi, K., Rose, G. C., and Mahan, C. J.: Ultrastructure of the dento-epithelial junction. J Periodont Res 11: 313, 1976.
244. Jones, S. J.: The tooth surface in periodontal disease. Dent Pract 22: 462, 1972.
245. Box, H. K.: Twelve Periodontal Studies. Toronto, Univ. Toronto Press, 1940.
246. Schroeder, H. E., and Listgarten, M. A.: Fine Structure of the Developing Epithelial Attachment of Human Teeth, p 66. Basel, S. Karger, 1971.
247. Hodson, J. J.: Origin and nature of the cuticula dentis. Nature 209: 990, 1966.
248. Stallard, R. E., Diab, M. A., and Zander, H. A.: The attaching substance between enamel and epithelium--a product of the epithelial cells. J Periodontol 36: 130, 1965.
249. Wertheimer, F. W., and Fullmer, H. M.: Morphologic and histochemical observations on the human dental cuticle. J Periodontol 33: 29, 1962.
250. Kobayashi, K., and Rose, G. G.: Ultrastructural histochemistry of the dento-epithelial junction. 3. Chloramine T-silver methenamine. J Periodont Res 14: 123, 1979.
251. Toto, P. D., and Sicher, H.: The epithelial attachment. Periodontics 2: 154, 1964.

252. Toto, P. D., and Sicher, H.: Mucopolysaccharides in the epithelial attachment. J Dent Res 44: 451, 1965.
253. Kobayashi, K., Rose, G. G., and Mahan, C. J.: Ultrastructural histochemistry of the dento-epithelial junction. I. Phosphotungstic acid and periodic acid-silver methenamine and periodic acid-thiosemicarbazide-silver proteinate. J Periodont Res 12: 351, 1977.
254. Kobayashi, K., and Rose, G. G.: Ultrastructural histochemistry of the dento-epithelial junction. II. Colloidal thorium and ruthenium red. J Periodont Res 13: 164, 1978.
255. Buchanan, W., and Bernard, G. W.: A technique for the removal of junctional epithelium from the epithelial attachment. J Dent Res 57: 895, 1978.
256. Waerhaug, J.: Enamel cuticle. J Dent Res 35: 313, 1956.
257. Cimasoni, G., Fiore-Donno, G., and Held, A. J.: Mucopolysaccharides in human epithelial reattachment. Helv Odont Acta 7: 60, 1963.
258. Frank, R., Fiore-Donno, G., Cimasoni, G., and Ogilvie, A.: Gingival reattachment after surgery in man: An electron microscopic study. J Periodontol 43: 597, 1972.
259. McHugh, W. D.: Some aspects of the development of gingival epithelium. Periodontics 1: 239, 1963.
260. Taylor, A. C., and Campbell, M. M.: Reattachment of gingival epithelium to the tooth. J Periodontol 43: 281, 1972.
261. Engler, W. O., Ramfjord, S. P., and Hiniker, J. J.: Development of epithelial attachment and gingival sulcus in rhesus monkeys. J Periodontol 36: 44, 1965.
262. Braga, A. M., and Squier, C. A.: Ultrastructure of regenerating junctional epithelium in the monkey. J Periodontol 51: 386, 1980.
263. Goldman, H. M.: Histologic topographic changes of inflammatory origin in the gingival fibers. J Periodontol 23: 104, 1952.
264. Gargiulo, A. W., Wentz, F. M., and Orban, B.: Dimensions and relations of the dentogingival junction in humans. J Periodontol 32: 261, 1961.
265. Selvig, K. A.: Nonbanded fibrils of collagenous nature in human periodontal connective tissue. J Periodont Res 3: 169, 1968.
266. Saglie, R., Johansen, J. R., and Flotra, L.: The zone of completely and partially destructed periodontal fibers in pathological pockets. J Clin Periodontol 2: 198, 1975.

267. Deporter, D. A., and Brown, D. Y.: Fine structural observations on the mechanism of loss of attachment during experimental periodontal disease in the rat. J Periodont Res 15: 304, 1980.
268. Shackelford, J. M.: Scanning electron microscopy of the dog periodontium. J Periodont Res 6: 45, 1971.
269. Gottlieb, B.: Biology of the cementum. J Periodontol 13: 13, 1942.
270. Gottlieb, B.: Continuous deposition of cementum. JADA 30: 842, 1943.
271. Bernier, J. L.: Changes in bone and cementum in periodontal disease. J Periodontol 22: 29, 1951.
272. Gustafson, A. G., and Persson, P-A.: The relationship between the direction of Sharpey's fibers and the deposition of cementum. Odont Tidskr 65: 457, 1957.
273. Kraw, A. G., and Enlow, D. H.: Continuous attachment of the periodontal membrane. Am J Anat 120: 133, 1967.
274. Orban, B.: The epithelial network in the periodontal membrane. JADA 44: 632, 1952.
275. Orban, B.: Resorption and repair on the surface of the root. JADA 15: 1768, 1928.
276. Coolidge, E. D.: The reaction of cementum in the presence of injury and infection. JADA 18: 499, 1931.
277. Steadman, S. R.: Resumé of the literature on root resorption. Angle Orthodont 12: 28, 1942.
278. Held, A-J.: Cementogenesis and the normal and pathologic structure of cementum. Oral Surg 4: 53, 1951.
279. Henry, J. L., and Weinmann, J. P.: The pattern of resorption and repair of human cementum. JADA 42: 270, 1951.
280. Southam, J. C.: Clinical and histologic aspects of peripheral cervical resorption. J Periodontol 38: 534, 1967.
281. Orban, B.: Gingivectomy or flap operation? JADA 26: 1276, 1939.
282. Borden, S. M.: Histologic study of healing following detachment of tissue as is commonly carried out in the vertical incision for the surgical removal of teeth. Canad Dent A J 14: 510, 1948.
283. Levine, H. L.: Periodontal flap surgery with gingival fiber retention. J Periodontol 43: 91, 1972.

284. Dedolph, T. H., and Clark, H. B.: A histological study of mucoperiosteal flap healing. J Oral Surg 16: 367, 1958.
285. Persson, P-A.: The regeneration of the marginal periodontium after flap operation. Acta Odont Scand 20: 43, 1962.
286. Levine, H. L., and Stahl, S. S.: Repair following periodontal flap surgery with the retention of gingival fibers. J Periodontol 43: 99, 1972.
287. Tonna, E. A., Stahl, S. S., and Asiedu, S.: A study of the reformation of severed gingival fibers in aging mice using ³H-proline autoradiography. J Periodont Res 15: 43, 1980.
288. Skillen, W. G., and Lundquist, G. R.: An experimental study of periodental membrane reattachment in healthy and pathologic tissues. JADA 24: 175, 1937.
289. Linghorne, W. J., and O'Connell, D. C.: Studies in the regeneration and attachment of supporting structures of the teeth. 1. Soft tissue reattachment. J Dent Res 29: 419, 1950.
290. Morris, M. L.: Healing of human periodontal tissues following surgical detachment from vital teeth: The position of the epithelial attachment. J Periodontol 32: 108, 1961.
291. Fish, E. W.: Parodontal disease: The pathology and treatment of chronic gingivitis. Brit Dent J 58: 531, 1935.
292. Jansen, M. T., Coppes, L., and Verdenus, H. H. W.: The healing of periodontal wounds in dogs. J Periodontol 26: 292, 1955.
293. Morris, M. L.: Healing of human periodontal tissues following surgical detachment: The arrangement of the fibers of the periodontal space. Periodontics 1: 118, 1963.
294. Nyman, S., and Karring, T.: Regeneration of surgically removed buccal alveolar bone in dogs. J Periodont Res 14: 86, 1979.
295. Ramfjord, S. P., Engler, W. O., and Hiniker, J. J.: A radioautographic study of healing following simple gingivectomy. II. The connective tissue. J Periodontol 37: 179, 1966.
296. Melcher, A. H.: Repair of wounds in the periodontium of the rat. Influence of periodontal ligament on osteogenesis. Archs Oral Biol 15: 1183, 1970.
297. Stahl, S. S.: Healing following simulated fiber retention procedures in rats. J Periodontol 48: 67, 1977.

298. Lahiffe, B. J., Caffesse, R. G., and Nasjleti, C. E.: Healing of periodontal flaps following use of MBR4197 (Flucrylate) in rhesus monkeys. J Periodontol 49: 635, 1978.
299. Caffesse, R. G., Ramfjord, S. P., and Nasjleti, C. E.: Reverse bevel periodontal flaps in monkeys. J Periodontol 39: 219, 1968.
300. Linghorne, W. J., and O'Connell, D. C.: Studies in the regeneration and reattachment of supporting structures of the teeth. II. Regeneration of alveolar process. J Dent Res 30: 604, 1951.
301. Melcher, A. H.: On the repair potential of periodontal tissues. J Periodontol 47: 256, 1976.
302. Karring, T., Cumming, B. R., Oliver, R. C., and Löe, H.: The origin of granulation tissue and its impact on postoperative results of mucogingival surgery. J Periodontol 46: 577, 1975.
303. Gould, T. R. L., Melcher, A. H., and Brunette, D. M.: Migration and division of progenitor cell populations in periodontal ligament after wounding. J Periodont Res 15: 20, 1980.
304. Line, S. E., Polson, A. M., and Zander, H. A.: Relationship between periodontal surgery, selective cell repopulation and ankylosis. J Periodontol 45: 725, 1974.
305. Karring, T., Nyman, S., and Lindhe, J.: Healing following implantation of periodontitis affected roots into bone tissues. J Clin Periodontol 7: 96, 1980.
306. Löe, H., and Waerhaug, J.: Experimental replantation of teeth in dogs and monkeys. Archs Oral Biol 3: 176, 1961.
307. Nyman, S., Karring, T., Lindhe, J., and Plantén, S.: Healing following implantation of periodontitis-affected roots into gingival connective tissue. J Clin Periodontol 7: 394, 1980.
308. Adell, R.: Regeneration of the periodontium. Scand J Plast Reconstr Surg Suppl 11, 1974.
309. Morris, M. L.: Healing of human periodontal tissues following surgical detachment and extirpation of vital pulps. J Periodontol 31: 23, 1960.
310. Morris, M. L.: The reattachment of human periodontal tissues following surgical detachment: A clinical and histological study. J Periodontol 24: 220, 1953.
311. Morris, M. L., and Thompson, R. H.: Healing of human periodontal tissues following surgical detachment: Factors related to the deposition of new cementum on dentin. Periodontics 1: 189, 1963.

312. Edwards, R. C., Ratcliff, P. A., and Raust, G. T.: Repair of the dentogingival interface. Dent Pract 19: 301, 1969.
313. Nalbandian, J., and Frank, R. M.: Electron microscopic study of the regeneration of cementum and periodontal connective tissue attachment in the cat. J Periodont Res 15: 71, 1980.
314. Morris, M. L.: Healing of human periodontal tissues following surgical detachment from non-vital teeth. J Periodontol 28: 222, 1957.
315. Erausquin, J., and Muruzábal, M.: Necrosis of cementum induced by root canal treatments in the molar teeth of rats. Archs Oral Biol 12: 1123, 1967.
316. Diem, C. R., Bowers, G. M., Ferrigno, P. D., and Fedi, P. F.: Regeneration of the attachment apparatus on pulpless teeth denuded of cementum in the rhesus monkey. J Periodontol 45: 18, 1974.
317. Helldén, L.: Periodontal healing following experimental injury to root surfaces of human teeth. Scand J Dent Res 80: 197, 1972.
318. Hill, T. J.: Regeneration of periodontal membrane after root curetment. Dent Cosmos 73: 799, 1931.
319. Bodecker, C. F.: Concerning the re-attachment of the periodontal membrane to cementum. J Dent Educ 6: 236, 1942.
320. Sorrin, S.: Root therapy and periodontal treatment in a case of advanced periodontal disease. J Dent Med 12: 130, 1957.
321. Hiatt, W. H.: Regeneration of the periodontium after endodontic therapy and flap operation. Oral Surg 12: 1471, 1959.
322. Rossman, S. R., Kaplowitz, B., and Baldinger, S. R.: Therapy of the endodontically and periodontally involved tooth. Oral Surg 13: 361, 1960.
323. Hiatt, W. H.: Regeneration via flap operation and the pulpal periodontal lesion. Periodontics 4: 205, 1966.
324. Register, A. A.: Induced reattachment in periodontic-endodontic lesions by root demineralization in situ. Oral Surg 45: 774, 1978.
325. Moskow, B. S.: The response of the gingival sulcus to instrumentation: A histologic investigation. I. The scaling procedure. J Periodontol 33: 282, 1962.
326. O'Bannon, J. Y.: The gingival tissues before and after scaling the teeth. J Periodontol 35: 69, 1964.
327. Schaffer, E. M., Stende, G., and King, D.: Healing of periodontal pocket tissues following ultrasonic scaling and hand planing. J Periodontol 35: 140, 1964.
328. Waerhaug, J.: Microscopic demonstration of tissue reaction incident to removal of subgingival calculus. J Periodontol 26: 26, 1955.
329. Kon, S., Novaes, A. B., Ruben, M. P., and Goldman, H. M.: Visualization of microvascularization of the healing periodontal wound. II. Curettage. J Periodontol 40: 96, 1969.

330. Lopez, N. J., and Belvederessi, M.: Subgingival scaling with root planing and curettage: Effects upon gingival inflammation. A Comparative study. J Periodontol 48: 354, 1977.
331. Moskow, B. S.: The response of the gingival sulcus to instrumentation: A histologic investigation. 2. Gingival Curettage. J Periodontol 35: 112, 1964.
332. Goldman, H. M.: Curettage by ultrasonic instrument. Oral Surg 13: 43, 1960.
333. Nadler, H.: Removal of crevicular epithelium by ultrasonic cures. J Periodontol 33: 220, 1962.
334. Stahl, S. S., Weiner, J. M., Benjamin, S., and Yamada, L.: Soft tissue healing following curettage and root planing. J Periodontol 42: 678, 1971.
335. Sanderson, A. D.: Gingival curettage by hand and ultrasonic instruments: A histologic comparison. J Periodontol 37: 279, 1966.
336. Morris, M. L.: The removal of pocket and attachment epithelium in humans: A histological study. J Periodontol 25: 7, 1954.
337. Raust, G. T.: What is the value of gingival curettage in periodontal therapy? Periodont Abstr 17: 142, 1969.
338. Carranza, F. A.: A technique for treating infrabony pockets so as to obtain reattachment. Dent Clin N Am p. 75, March 1960.
339. Gibbins, J. R.: Migration of stratified squamous epithelium in vivo. Am J Path 53: 929, 1968.
340. Sato, M.: Histopathological study of healing processes after surgical treatment for alveolar pyorrhea. Tokyo Dent Col Bull Oral Path 2: 45, 1958.
341. Sato, M.: Histopathological study of the healing process after surgical treatment for alveolar pyorrhea. Tokyo Dent Col Bull 1: 71, 1960.
342. Blass, J. L., and Lite, T.: Gingival healing following surgical curettage: A histopathologic study. N Y S Dent J 25: 127, 1959.
343. Goldman, H. M.: Histologic assay of healing following ultrasonic curettage versus hand-instrumentation curettage. Oral Surg 14: 925, 1961.
344. Ingle, J. I.: Periodontal curettement in the premaxilla. J Periodontol 23: 143, 1952.
345. Shapiro, M.: Reattachment in periodontal disease. J Periodontol 24: 26, 1953.
346. Chaikin, B. S.: Subgingival curettage. J Periodontol 25: 240, 1954.
347. Everett, F. G.: Re-attachment in the treatment of periodontoclasia. Ore S Dent J 23: 2, (January) 1954.

348. Morris, M. L.: Healing of naturally occurring periodontal pockets about vital human teeth. J Periodontol 26: 285, 1955.
349. Schaffer, E. M.: The new attachment operation in subcrestal pockets. Oral Surg 11: 253, 1958.
350. Box, H. K.: Studies in periodontal pathology. Bull No. 7 Canad Dent Res Found, May 1924.
351. Schaffer, E. M., and Zander, H. A.: Histologic evidence of reattachment of periodontal pockets. Paradentologie 7: 101, 1953.
352. Cross, W. G.: Reattachment following curettage. Dent Pract 7: 38, 1956.
353. McCall, J. O.: Tissue regeneration in periodontal disease. J Periodontol 34: 47, 1963.
354. Rule, R. W.: Reattachment. Case report. J Periodontol 14: 8, 1943.
355. Benjamin, E. M.: The quantitative comparison of subgingival curettage and gingivectomy in the treatment of periodontitis simplex. J Periodontol 27: 144, 1956.
356. Rateitschak, K. H.: The therapeutic effect of local treatment on periodontal disease assessed upon evaluation of different diagnostic criteria. J Periodontol 35: 155, 1964.
357. Wade, A. B.: Causes of failure in treatment of localized pocketing. Dent Health 4: 9, 1965.
358. Zamet, J. S.: A comparative clinical study of three periodontal surgical techniques. J Clin Periodontol 2: 87, 1975.
359. Korn, N. A., Schaffer, E. M., and McHugh, R. B.: An experimental assessment of gingivectomy and soft tissue curettage in dogs. J Periodontol 36: 96, 1965.
360. Burgett, F. G., Knowles, J. W., Nissle, R. R., Shick, R. A., and Ramfjord, S. P.: Short term results of three modalities of periodontal treatment. J Periodontol 48: 131, 1977.
361. Knowles, J. W., Burgett, F. G., Nissle, R. R., Shick, R. A., Morrison, E. C., and Ramfjord, S. P.: Results of periodontal treatment related to pocket depth and attachment level. Eight years. J Periodontol 50: 225, 1979.
362. McCall, J. O.: An improved method of inducing reattachment of the gingival tissues in periodontoclasia. Dent Items Interest 48: 342, 1926.

363. Miller, S. C., and Sorrin, S.: The action and use of sodium sulfid solution as an epithelial solvent. Dent Cosmos 69: 1113, 1927.
364. Bjorndahl, O.: The conservative treatment procedure of the complex pocket. J Periodontol 11: 31, 1940.
365. Beube, F. E.: An experimental study of the use of sodium sulphid solution in treatment of periodontal pockets. J Periodontol 10: 49, 1939.
366. Kaplan, H., and Mann, J. B.: How is pyorrhea cured? A comparative study of several methods for treatment of pyorrhea alveolaris. JADA 29: 1471, 1942.
367. Glickman, I., and Patur, B.: Histologic study of the effects of antiformin on the soft tissue wall of periodontal pockets in human beings. JADA 51: 420, 1955.
368. Johnson, R. E., and Waerhaug, J.: Effect of antiformin on gingival tissues. J Periodontol 27: 24, 1956.
369. Løe, H.: Chemical gingivectomy, effect of potassium hydroxide on periodontal tissues. Acta Odont Scand 19: 517, 1961.
370. Younger, W. J.: Some of the latest phases in implantation and other operations. Dent Cosmos 35: 102, 1893.
371. Barkann, L.: A conservative surgical technic for the eradication of a pyorrhea pocket. JADA 26: 61, 1939.
372. Rosenthal, S. L.: Conservative methods for reduction of the periodontal pocket. Dent Items Interest 72: 154, 1950.
373. Ramfjord, S. P.: Reattachment in periodontal therapy. JADA 45: 513, 1952.
374. Waerhaug, J., and Løe, H.: Effect of phenol camphor on gingival tissue. J Periodontol 29: 59, 1958.
375. Linghorne, W. J., and O'Connell, D. C.: Studies in the reattachment and regeneration of the supporting structures of the teeth. III. Regeneration in epithelized pockets. J Dent Res 34: 164, 1955.
376. Linghorne, W. J.: Studies in the reattachment and regeneration of the supporting structures of the teeth. IV. Regeneration in epithelized pockets following organization of a blood clot. J Dent Res 36: 4, 1957.
377. Berliner, A.: Elimination of periodontal pockets. Dent Digest 56: 397, 1950.

378. Wertheimer, F. W.: Effectiveness of "Berliner Epithelial Scalpel" in removing the epithelial lining of periodontal pockets. J Periodontol 25: 264, 1954.
379. Ramfjord, S. P., and Ash, M. M. Jr.: Periodontology and Periodontics, p. 603. Philadelphia, W. B. Saunders, 1979.
380. Ramfjord, S. P.: Present status of the modified Widman flap procedure. J Periodontol 48: 558, 1977.
381. Ewen, S. J.: Ultrasonic surgery in periodontal therapy. N Y S Dent J 25: 189, 1959.
382. Zach, L.: The histology of the response to ultrasonic curettage. J Dent Res 40: 751, 1961. IADR Abstr No. 305.
383. Ewen, S. J.: The ultrasonic wound -- some microscopic observations. J Periodontol 32: 315, 1961.
384. Frisch, J., Bhaskar, S., and Shell, D.: Effect of ultrasonic instrumentation on human gingival connective tissue. Periodontics 5: 123, 1967.
385. Ramfjord, S. P., and Ash, M. M. Jr.: Periodontology and Periodontics, p. 523. Philadelphia, W. B. Saunders, 1979.
386. Hagerman, D. A., and Arnim, S. S.: The effect of periodontal treatment on the connective tissue fibers of the marginal gingiva. JADA 48: 158, 1954.
387. Glickman, I., Goldman, H. M., Thomas, B. O. A., and Ramfjord, S. P.: Is scaling prior to gingival surgery a necessary procedure? Periodont Abstr 4: 99, 1956.
388. Gottsegen, R.: Should teeth be scaled prior to surgery? J Periodontol 32: 301, 1961.
389. Ramfjord, S. P., and Nissle, R. R.: The modified Widman flap. J Periodontol 45: 601, 1974.
390. Ramfjord, S. P., and Ash, M. M. Jr.: Periodontology and Periodontics, p. 500. Philadelphia, W. B. Saunders, 1979.
391. Grillo, H. C.: Derivation of fibroblasts in the healing wound. Arch Surg 88: 218, 1964.
392. Collings, C. K.: The role of the blood clot in periodontal treatment. J Periodontol 22: 171, 1951.
393. Garant, P. R., and Cho, M.: Cytoplasmic polarization of periodontal ligament fibers. J Periodont Res 14: 95, 1979.

394. Stahl, S. S., Slavkin, H. C., Yamada, L., and Levine, S.: Speculations about gingival repair. J Periodontol 43: 395, 1972.
395. Friedman, N.: Reattachment and roentgenograms. J Periodontol 29: 98, 1958.
396. Yukna, R. A.: A clinical and histologic study of healing following the excisional new attachment procedure in rhesus monkeys. J Periodontol 47: 701, 1976.
397. Cook, R. T., Hutchens, L. H., and Burks, E. J.: Periodontal osseous defects associated with vitally submerged roots. J Periodontol 48: 249, 1977.
398. Caton, J., Nyman, S., and Zander, H.: Histometric evaluation of periodontal surgery. II. Connective tissue attachment levels after four regenerative procedures. J Clin Periodontol 7: 224, 1980.
399. Beube, F. E.: Interdental tissue resection: An experimental study of a surgical technique which aids in repair of the periodontal tissues to their original contour and function. Am J Ortho Oral Surg 33: 497, 1947.
400. Morris, M. L.: The unrepositioned muco-periosteal flap. Periodontics 3: 147, 1965.
401. Raust, G. T.: Mini-flaps. J Periodontol 40: 56, 1969.
402. Klavan, B.: The replaced graft. J Periodontol 41: 406, 1970.
403. Sandalli, P., and Wade, A. B.: Alterations in crevicular fluid flow during healing following gingivectomy and flap procedures. J Periodont Res 4: 314, 1969.
404. Barletta, B. O., Caffesse, R. G., Paladino, C., and Plot, C. R.: Comparative biometric evaluation of results obtained after gingivectomy and reverse-bevel periodontal flap surgery. J Dent Res 51: 1227, 1972. Argent. Sect., IADR Abstr No. 33.
405. Wade, A. B.: An assessment of the flap operation. Dent Pract 13: 11, 1962.
406. Yukna, R. A., Bowers, G. M., Lawrence, J. J., and Fedi, P. F.: A clinical study of healing in humans following the excisional new attachment procedure. J Periodontol 47: 696, 1976.
407. Yukna, R. A.: Longitudinal evaluation of the excisional new attachment procedure in humans. J Periodontol 49: 142, 1978.
408. Yukna, R. A., and Williams, J. E.: Five year evaluation of the excisional new attachment procedure. J Periodontol 51: 382, 1980.

409. Smith, D. H., Ammons, W. F., and Van Belle, G.: A longitudinal study of periodontal status comparing osseous recontouring with flap curettage. I. Results after 6 months. J Periodontol 51: 367, 1980.
410. Sullivan, H. C., and Atkins, J. H.: Free autogenous gingival grafts. III. Utilization of grafts in the treatment of gingival recession. Periodontics 6: 152, 1968.
411. Mlinek, A., Smuckler, H., and Buchner, A.: The use of free gingival grafts for the coverage of denuded roots. J Periodontol 44: 248, 1973.
412. Livingston, H. L.: Total coverage of multiple and adjacent root surfaces with a free gingival autograft. J Periodontol 46: 209, 1975.
413. Matter, J., and Cimasoni, G.: Creeping attachment after free gingival grafts. J Periodontol 47: 574, 1976.
414. Grupe, H. E., and Warren, R. F.: Repair of gingival defects by a sliding flap operation. J Periodontol 27: 92, 1956.
415. Rosenberg, M. M.: A laterally repositioned flap involving a non-vital tooth: Case report. Periodontics 1: 164, 1963.
416. McFall, W. T.: The laterally repositioned flap -- criteria for success. Periodontics 5: 89, 1967.
417. Cohen, D. W., and Ross, S. E.: The double papilla repositioned flap in periodontal therapy. J Periodontol 39: 65, 1968.
418. Björn, H.: Coverage of denuded root surfaces with a lateral sliding flap. Use of free gingival grafts. Odont Revy 22: 37, 1971.
419. Smuckler, H.: Lateral positioned mucoperiosteal pedicle grafts in the treatment of denuded roots. J Periodontol 47: 590, 1976.
420. Guinard, E. A., and Caffesse, R. G.: Treatment of localized gingival recessions. Part I. Lateral sliding flap. J Periodontol 49: 351, 1978.
421. Caffesse, R. G., and Guinard, E. A.: Treatment of localized gingival recessions. Part IV. Results after three years. J Periodontol 51: 167, 1980.
422. Wilderman, M. N., and Wentz, F. M.: Repair of a dentogingival defect with a pedicle flap. J Periodontol 36: 218, 1965.
423. Sugarman, E. F.: A clinical and histological study of the attachment of grafted tissue to bone and teeth. J Periodontol 40: 381, 1969.
424. Pfeiffer, J. S., and Heller, R.: Histologic evaluation of full and partial thickness lateral repositioned flaps: A pilot study. J Periodontol 42: 331, 1971.

425. Goldman, H. M., and Shuman, A.: A technic for the therapy of the two-walled osseous defect. Periodontics 6: 88, 1968.
426. Goldman, H. M., Shuman, A. M., and Isenberg, G. A.: Management of the partial furcation involvement. Periodontics 6: 197, 1968.
427. Hiatt, W. H.: The repositioned alveolar ridge mucosal flap. Periodontics 5: 132, 1967.
428. Ruben, M. P., Goldman, H. M., and Janson, W.: Biologic considerations fundamental to successful employment of laterally repositioned pedicle flaps and free autogenous gingival grafts in periodontal therapy. In Stahl, S. S. (ed.) Periodontal Surgery Biologic Basis and Technique, p. 235. Springfield, C. C. Thomas, 1976.
429. Simpson, H. E.: The reattachment of mucoperiosteal flaps in surgical extraction wounds in *Macacus rhesus* monkeys. Austral Dent J 4: 86, 1959.
430. Melcher, A. H.: Wound healing in monkey (*Macaca irus*) mandible: Effect of elevating periosteum on formation of subperiosteal callus. Archs Oral Biol 16: 461, 1971.
431. Melcher, A. H., and Accursi, G. E.: Osteogenic capacity of periosteal and osteoperiosteal flaps elevated from the parietal bone of the rat. Archs Oral Biol 16: 573, 1971.
432. Beasley, J. D., Hunsuck, E. E., Bhaskar, S. N., and Cutright, D. E.: Autogenous osteomucoperiosteal pedicle grafts in miniature swine. J Periodontol 43: 146, 1972.
433. Goldman, H. M., and Smukler, H.: Controlled surgical stimulation of periosteum. J Periodontol 49: 518, 1978.
434. Smukler, H., and Goldman, H. M.: Laterally repositioned "stimulated" osteoperiosteal pedicle grafts in the treatment of denuded roots. J Periodontol 50: 379, 1979.
435. Passanezi, E., Alves, M. E. A. F., Janson, W. A., and Ruben, M. P.: Periosteal activation and root demineralization associated with the horizontal sliding flap. J Periodontol 50: 384, 1979.
436. Krømer, H.: Bone homografts in minor oral surgery. Proc Roy Soc Med 55: 607, 1962.
437. Lockhart, R. L.: Free periosteal autografts in the treatment of periodontal defects. Thesis, Univ. Missouri/Kansas City, 1975.
438. Hegedus, Z.: The rebuilding of the alveolar process by bone transplantation. Dent Cosmos 65: 736, 1923.

439. Norberg, O.: Biologi och terapi. Svensk Tandl Tidskr 34: 32, 1941.
440. Nordenram, A., and Landt, H.: Evaluation of a surgical technique in the periodontal treatment of maxillary anterior teeth. Acta Odont Scand 27: 283, 1969.
441. Harvey, P. M.: Management of advanced periodontitis. Part I. Preliminary report of a method of surgical reconstruction. New Zeal Dent J 61: 180, 1965.
442. Restrepo, O. J.: Coronally repositioned flap: Report of four cases. J Periodontol 44: 564, 1973.
443. Brustein, D. D.: Cosmetic dentistry/coronally repositioned pedicle graft. Dent Survey 46: 22, (July) 1970.
444. Maynard, J. G.: Coronal positioning of a previously placed autogenous gingival graft. J Periodontol 48: 151, 1977.
445. Kirkham, D. B., and Hoge, H. W.: Repositioning free gingival grafts to cover denuded root surfaces. Dent Survey 56: 46, (July) 1980.
446. Bernimoulin, J. P., Lüscher, B., and Mühlemann, H. R.: Coronally repositioned periodontal flap. J Clin Periodontol 2: 1, 1975.
447. Guinard, E. A., and Caffesse, R. G.: Treatment of localized gingival recessions. Part III. Comparison of results obtained with lateral sliding and coronally repositioned flaps. J Periodontol 49: 457, 1978.
448. Kalmi, J., Moscor, M., and Goranov, Z.: The solution of the esthetic problem in the treatment of periodontal disease of anterior teeth: Gingivoplastic operation. Paradentologie 3: 53, 1949.
449. Patur, B., and Glickman, I.: Gingival pedicle flaps for covering root surfaces denuded by chronic destructive periodontal disease -- a clinical experiment. J Periodontol 29: 50, 1958.
450. Sumner, C. F.: Surgical repair of recession on maxillary cuspid: Initially repositioning the gingival tissues. J Periodontol 40: 119, 1969.
451. Schluger, S., Yuodelis, R. A., and Page, R. C.: Periodontal Disease, p. 483. Philadelphia, Lea and Febiger, 1977.
452. Morris, M. L.: Healing of naturally occurring periodontal pockets: Preliminary report of the effects of external pressure. J Periodontol 32: 202, 1961.
453. Ramfjord, S. P., and Ash, M. M. Jr.: Periodontology and Periodontics, p. 531. Philadelphia, W. B. Saunders, 1979.
454. Ruben, M. P., Smuckler, H., Schulman, S. M., Kon, S., and Bloom, A. A.: Healing of periodontal surgical wounds. In Goldman, H. M., and Cohen, D. W., Periodontal Therapy, 6th ed., p. 640. St. Louis, C. V. Mosby, 1980.

455. Baum, B. J., and Wright, W. E.: Demonstration of fibronectin as a major extracellular protein of human gingival fibroblasts. J Dent Res 59: 631, 1980.
456. Yajima, T., Rose, G. R., and Mahan, C. J.: Human gingival fibroblast cell lines in vitro. II. Electron microscopic studies of fibrogenesis. J Periodont Res 15: 267, 1980.
457. Boyko, G. A., Brunette, D. M., and Melcher, A. H.: Cell attachment to demineralized root surfaces in vitro. J Periodont Res 15: 297, 1980.
458. Weiss, R. E., and Reddi, A. H.: Synthesis and localization of fibronectin during collagenous matrix-mesenchymal cell interaction and differentiation of cartilage and bone in vivo. Proc Nat Acad Sci 77: 2074, 1980.
459. Lloyd, C.: Fibronectin: A function at the junction. Nature 279: 473, 1979.
460. Goldman, H. M.: Subgingival curettage, a rationale. J Periodontol 19: 54, 1948.
461. Goldman, H. M., and Cohen, D. W.: The infrabony pocket: Classification and treatment. J Periodontol 29: 272, 1958.
462. Melcher, A. H.: Healing of wounds in the periodontium. In Melcher, A. H., and Bowen, W. H., (eds.), Biology of the Periodontium, p. 509. New York, Academic Press, 1969.
463. Adams, D. F.: Implant therapy of infrabony pockets. Periodont Abstr 14: 63, 1966.
464. Nabers, C. L., and O'Leary, T. J.: Autogenous bone grafts: Case Report. Periodontics 5: 251, 1967.
465. Fullmer, H. M.: A histochemical study of periodontal disease in the maxillary alveolar processes of 135 autopsies. J Periodontol 32: 206, 1961.
466. Carranza, F. A., and Glickman, I.: Some observations on the microscopic features of infrabony pockets. J Periodontol 28: 33, 1957.
467. Goldman, H. M.: The behavior of transseptal fibers in periodontal disease. J Dent Res 36: 249, 1957.
468. Prichard, J. F.: The etiology, diagnosis and treatment of the intrabony defect. J Periodontol 38: 455, 1967.
469. Toto, P. D., Pollack, R. J., and Gargiulo, A. W.: Pathogenesis of periodontitis. Periodontics 2: 197, 1964.
470. Bell, D. G.: A case of reattachment? J Periodontol 8: 30, 1937.
471. Bjorndahl, O.: Reattachment and bone regeneration: Report of case. JADA 36: 356, 1948.

472. McCall, J. O.: Periodontal reattachment: Clinical and roentgenological evidence. JADA 43: 382, 1951.
473. Cahn, L. R.: Hyperemic decalcification. Oral Surg 7: 332, 1954.
474. Prichard, J.: Regeneration of bone following periodontal therapy. Oral Surg 10: 247, 1957.
475. Goldstein, A.: Bone regeneration with conservative periodontal treatment. J Dent Med 13: 109, 1958.
476. Nabers, J. M., Meador, H. L., Nabers, C. L., and O'Leary, T. J.: Chronology, an important factor in the repair of osseous defects. Periodontics 2: 304, 1964.
477. Moskow, B. S., Wasserman, B. H., Hirschfeld, L. S., and Morris, M. L.: Repair of periodontal tissues following acute localized osteomyelitis. Periodontics 5: 29, 1967.
478. Bitter, N. C.: Periodontal infection treated conservatively by scaling. Oral Surg 28: 484, 1969.
479. Fleming, W. C.: A clinical and microscopical study of periodontal tissue treated by instrumentation. Pacific Dent Gazette 34: 568, 1926.
480. Smith, T. S.: The treatment of two periodontal cases. J Periodontol 20: 129, 1949.
481. Goldman, H. M.: A rationale for the treatment of the intrabony pocket; one method of treatment, subgingival curettage. J Periodontol 20: 83, 1949.
482. Prichard, J.: The infrabony pocket as a predictable procedure. J Periodontol 28: 202, 1957.
483. Schreiber, H. R.: Successful treatment of a deep two wall intrabony pocket. J Periodontol 32: 342, 1961.
484. Kramer, G. M., and Kohn, D. J.: Post-operative care of the infra-bony pocket. J Periodontol 32: 95, 1961.
485. Prichard, J.: Gingivoplasty, gingivectomy, and osseous surgery. J Periodontol 32: 275, 1961.
486. Moskow, B. S.: Healing potential in periodontal lesions. The interdental crater. J Periodontol 48: 754, 1977.
487. Moskow, B. S.: Repair potential in periodontal disease: The interproximal channel. J Periodontol 49: 55, 1978.
488. Theilade, J.: An evaluation of the reliability of radiographs in the measurement of bone loss in periodontal disease. J Periodontol 31: 143, 1960.

489. Bender, I. B., and Seltzer, S.: Roentgenographic and direct observation of experimental lesions in bone: II. JADA 62: 708, 1961.
490. Burnette, E. W.: Limitation of the roentgenogram in periodontal diagnosis. J Periodontol 42: 293, 1971.
491. Goldman, H. M., and Stallard, R. E.: Limitations of the radiograph in the diagnosis of osseous defects in periodontal disease. J Periodontol 44: 626, 1973.
492. Patur, B., and Glickman, I.: Clinical and roentgenographic evaluation of the post treatment healing of infrabony pockets. J Periodontol 33: 164, 1962.
493. Ellegaard, B., and Løe, H.: New attachment of periodontal tissues after treatment of intrabony lesions. J Periodontol 42: 648, 1971.
494. Patur, B.: Osseous defects: Evaluation of diagnostic and treatment methods. J Periodontol 45: 523, 1974.
495. Polson, A. M., and Heijl, L. C.: Osseous repair in infrabony periodontal defects. J Clin Periodontol 5: 13, 1978.
496. Rosling, B., Nyman, S., and Lindhe, J.: The effect of systemic plaque control on bone regeneration in infrabony pockets. J Clin Periodontol 3: 38, 1976.
497. Wade, A. B.: The flap operation. J Periodontol 37: 95, 1966.
498. Rosling, B., Nyman, S., Lindhe, J., and Barbro, J.: The healing potential of the periodontal tissues following different techniques of periodontal surgery in plaque-free dentitions. J Clin Periodontol 3: 233, 1976.
499. Caton, J., and Zander, H. A.: Osseous repair of an infrabony pocket without new attachment of connective tissue. J Clin Periodontol 3: 54, 1976.
500. Caton, J., and Nyman, S.: Histometric evaluation of periodontal surgery. I. The modified Widman flap procedure. J Clin Periodontol 7: 212, 1980.
501. Ellegaard, B., Karring, T., Davies, R., and Løe, H.: New attachment after treatment of intrabony defects in monkeys. J Periodontol 45: 368, 1974.
502. Rockoff, S. C., Rockoff, H. S., and Sackler, A. M.: Reattachment -- A case in point. J Periodontol 29: 261, 1958.
503. Hiatt, W. H., Schallhorn, R. G., and Aaronian, A. J.: The induction of new bone and cementum formation. IV. Microscopic examination of the periodontium following human bone and marrow allograft, autograft and nongraft periodontal regenerative procedures. J Periodontol 49: 495, 1978.

504. Listgarten, M. A., and Rosenberg, M. M.: Histological study of repair following new attachment procedures in human periodontal lesions. J Periodontol 50: 333, 1979.
505. Bjorn, H.: Experimental studies on reattachment. Dent Pract 11: 351, 1961.
506. Bjorn, H., Hollender, L., and Lindhe, J.: Tissue regeneration in patients with periodontal disease. Odont Revy 16: 315, 1965.
507. Ellegaard, B., Karring, T., and Løe, H.: New periodontal attachment procedure based on retardation of epithelial migration. J Clin Periodontol 1: 75, 1974.
508. Ellegaard, B., Karring, T., and Løe, H.: Retardation of epithelial migration in new attachment attempts in intrabony defects in monkeys. J Clin Periodontol 3: 23, 1976.
509. Sicher, H.: Some biologic aspects of bone grafting. Dent Clin N Am p. 427, July 1962.
510. Cross, W. G.: Bone implants in periodontal diseases -- a further study. J Periodontol 28: 184, 1957.
511. Mann, W. V.: Autogenous transplant in the treatment of an infrabony pocket: Case report. Periodontics 2: 205, 1964.
512. Ewen, S. J.: Bone swaging. J Periodontol 36: 57, 1965.
513. Nabers, C. L., and O'Leary, T. J.: Autogenous bone transplants in the treatment of osseous defects. J Periodontol 36: 5, 1965.
514. Ross, S. E., Malamed, E. H., and Amsterdam, M.: The contiguous autogenous transplant -- its rationale, indications and technique. Periodontics 4: 246, 1966.
515. Burnette, E. W.: Treatment of an isolated, severe osseous crater. J Periodontol 38: 148, 1967.
516. Robinson, R. E.: Osseous coagulum for bone induction. J Periodontol 40: 503, 1969.
517. Halliday, D. G.: The grafting of newly formed autogenous bone in the treatment of osseous defects. J Periodontol 40: 511, 1969.
518. Robinson, R. E.: The osseous coagulum for bone induction technique. Calif Dent A J 46: 18, 1970.
519. Rosenberg, M. M.: Free osseous tissue autografts as a predictable procedure. J Periodontol 42: 195, 1971.
520. Rosenberg, M. M.: Re-entry of an osseous defect treated by a bone implant after a long duration. J Periodontol 42: 360, 1971.

521. Soehren, S. E., and Van Swol, R. L.: The healing extraction site: A donor area for potential grafting material. J Periodontol 50: 128, 1979.
522. Rosenberg, E. S., Garber, D. A., and Abrams, B.: Repair of bony defect using an intraoral exostosis as the donor site. J Periodontol 50: 476, 1979.
523. Matsue, I., Collings, C. K., Zimmerman, E. R., and Vail, W. C.: Microdensitometric analysis of human autogenous alveolar bone implants. J Periodontol 41: 489, 1970.
524. Matsue, I., Zimmerman, E. L., Collings, C. K., and Best, J. T.: Microdensitometric analysis of human autogenous bone implants. II. Two dimensional density and pattern analysis of interproximal alveolar bone. J Periodontol 42: 435, 1971.
525. Hiatt, W. H., and Schallhorn, R. G.: Intraoral transplants of cancellous bone and marrow in periodontal lesions. J Periodontol 44: 194, 1973.
526. Froum, S. J., Thaler, R., Scopp, I. W., and Stahl, S. S.: Osseous autografts. I. Clinical responses to bone blend or hip marrow grafts. J Periodontol 46: 515, 1975.
527. Froum, S. J., Ortiz, M., Witkin, R. T., Thaler, R., Scopp, I. W. and Stahl, S. S.: Osseous autografts. III. Comparison of osseous coagulum-bone blend implants with open curettage. J Periodontol 47: 287, 1976.
528. Carraro, J. J., Sznajder, N., and Alonso, C. A.: Intraoral cancellous bone autografts in the treatment of infrabony pockets. J Clin Periodontol 3: 104, 1976.
529. Rivault, A. F., Toto, P. D., Levy, S., and Gargiulo, A.: Autogenous bone grafts: Osseous coagulum and osseous retrograde procedures in primates. J Periodontol 42: 787, 1971.
530. Yuktanandana, I.: Bone graft in the treatment of intrabony periodontal pockets in dogs. A histological investigation. J Periodontol 30: 17, 1959.
531. Patterson, R. L. Jr., Collings, C. K., and Zimmerman, E. R.: Autogenous implants in the alveolar process of the dog with induced periodontitis. Periodontics 5: 19, 1967.
532. Coverly, L., Toto, P., and Gargiulo, A.: Osseous coagulum: A histologic evaluation. J Periodontol 46: 596, 1975.
533. Ross, S. E., and Cohen, D. W.: The fate of a free osseous tissue autograft: A clinical and histologic case report. Periodontics 6: 145, 1968.
534. Nabers, C. L., Reed, O. M., and Hamner, J. E.: Gross and histologic evaluation of an autogenous bone graft 57 months postoperatively. J Periodontol 43: 702, 1972.

535. Hawley, C. E., and Miller, J.: A histologic examination of a free osseous autograft. J Periodontol 46: 289, 1975.
536. Froum, S. J., Thaler, R., Scopp, I. W., and Stahl, S. S.: Osseous autografts. II. Histologic responses to osseous coagulum-bone blend grafts. J Periodontol 46: 656, 1975.
537. Moskow, B. S., Karsh, F., and Stein, S. D.: Histologic assessment of autogenous bone graft. J Periodontol 50: 291, 1979.
538. Forsberg, H.: Transplantation of os purum and bone chips in the surgical treatment of periodontal disease. Acta Odont Scand 13: 235, 1956.
539. Schallhorn, R. G.: Eradication of bifurcation defects utilizing frozen autogenous hip marrow implants. Periodont Abstr 15: 101, 1967.
540. Schallhorn, R. G.: The use of autogenous hip marrow biopsy implants for bony crater defects. J Periodontol 39: 145, 1968.
541. DeMarco, T. J., and Scaletta, L. J.: The use of autogenous hip marrow in the treatment of juvenile periodontosis: A case report. J Periodontol 41: 683, 1970.
542. Seibert, J. S.: Reconstructive periodontal surgery: Case Report. J Periodontol 41: 113, 1970.
543. Haggerty, P. C., and Maeda, I.: Autogenous bone grafts: A revolution in the treatment of vertical bone defects. J Periodontol 42: 626, 1971.
544. Schallhorn, R. G.: Postoperative problems associated with iliac transplants. J Periodontol 43: 3, 1972.
545. Burnette, E. W.: Fate of an iliac crest graft. J Periodontol 43: 88, 1972.
546. Dragoo, M. R., and Sullivan, H. C.: A clinical and histological evaluation of autogenous iliac bone grafts in humans. Part II. External root resorption. J Periodontol 44: 614, 1973.
547. Hoffman, I. D., and Flanagan, P.: Exophytic granulation reactions associated with autogenous iliac marrow. J Periodontol 45: 586, 1974.
548. Schallhorn, R. G., Hiatt, W. H., and Boyce, W.: Iliac transplants in periodontal therapy. J Periodontol 41: 566, 1970.
549. Stabholz, A., Brayer, L., Gedalia, I., Yosipovitch, Z., and Soskolne, W. A.: Effect of fluoride on autogenous iliac cancellous bone and marrow transplants in surgically created intrabony pockets in dogs. J Periodontol 48: 413, 1977.
550. Ellegaard, B., Nielsen, I. M., and Karring, T.: Composite jaw and iliac cancellous bone grafts in intrabony defects in monkeys. J Periodont Res 11: 299, 1976.

551. Cross, W. G.: Bone-grafts in periodontal disease. Dent Pract 6: 98, 1955.
552. Krømer, H.: Surgical treatment of periodontal disease. Internat Dent J 8: 347, 1958.
553. Krømer, H.: Implantation of bone homografts in periodontal pockets. Dent Clin N Am p. 471, July 1962.
554. Haggerty, P. C.: Human allografts -- the efficient therapeutic approach to the infrabony defect. J Periodontol 48: 743, 1977.
555. Schallhorn, R. G., and Hiatt, W. H.: Human allografts of iliac cancellous bone and marrow in periodontal osseous defects. II. Clinical observations. J Periodontol 43: 67, 1972.
556. Strub, J. R., Garberthüel, T. W., and Firestone, A. R.: Comparison of tricalcium phosphate and frozen allogenic bone implants in man. J Periodontol 50: 624, 1979.
557. Mellonig, J. T., Bowers, G. M., Bright, R. W., and Lawrence, J. J.: Clinical evaluation of freeze-dried bone allografts in periodontal osseous defects. J Periodontol 47: 125, 1976.
558. Sepe, W. W., Bowers, G. M., Lawrence, J. J., Friedlander, G. E., and Koch, R. W.: Clinical evaluation of freeze-dried bone allografts in periodontal osseous defects -- Part II. J Periodontol 49: 9, 1978.
559. Altieri, E. T., Reeve, C. M., and Sheridan, P. J.: Lyophilized bone allografts in periodontal intraosseous defects. J Periodontol 50: 510, 1979.
560. Libin, B. M., Ward, H. L., and Fishman, L.: Decalcified, lyophilized bone allografts for use in human periodontal defects. J Periodontol 46: 51, 1975.
561. Hiatt, W. H.: The induction of new bone and cementum formation. III. Utilizing bone and marrow allografts in dogs. J Periodontol 41: 596, 1970.
562. Schreiber, H. R., Harder, E. W., and Thompson, L. D.: Cartilage and bone grafts in supra-bony pockets in dogs. J Periodontol 30: 291, 1959.
563. Cross, W. G.: The histology of a bone implant. Dent Pract 8: 26, 1957.
564. Hiatt, W. H., and Schallhorn, R. G.: Human allografts of iliac cancellous bone and marrow in periodontal osseous defects. I. Rationale and methodology. J Periodontol 42: 642, 1971.
565. Poulson, R. C., Rubenstein, A., and Garguilo, A. W.: Allogeneic iliac transplants in rhesus monkeys. J Periodontol 47: 187, 1976.

566. Hurt, W. C.: Freeze-dried bone homografts in periodontal lesions in dogs. J Periodontol 39: 89, 1968.
567. Narang, R., and Wells, H.: Bone induction in experimental periodontal bony defects in dogs with decalcified allogeneic bone matrix grafts. Oral Surg 33: 306, 1972.
568. Beube, F. E., and Silvers, H. F.: Further studies on bone generation with the use of boiled heterogenous bone. J Periodontol 7: 17, 1936.
569. Beube, F. E.: A study on reattachment of the supporting structures of the teeth. J Periodontol 18: 55, 1947.
570. Cross, W. G.: The use of bone implants in the treatment of periodontal pockets. Dent Clin N Am p. 107, March 1960.
571. Gurkan, S. I., and Melek, S.: Anorganic bone implantation in infrabony periodontal pockets. Internat Dent J 13: 703, 1963.
572. Older, L. B.: The use of heterogenous bovine bone implants in the treatment of periodontal pockets. J Periodontol 38: 539, 1967.
573. Shaw, J. G.: The treatment of multiple periodontal pockets by extended flap operation. Parodontologie 16: 121, 1962
574. Scopp, I. W., Morgan, F. H., Dooner, J. J., Fredrics, H. J., and Heyman, R. A.: Bovine bone (Boplant) implants for infrabony oral lesions. Periodontics 4: 169, 1966.
575. Beube, F. E., and Silvers, H. F.: Influence of devitalized heterogenous bone-powder on regeneration of alveolar and maxillary bone of dogs. J Dent Res 14: 15, 1934.
576. Beube, F. E.: Factors in the repair of alveolar bone and cementum Oral Surg 2: 379, 1949.
577. Beube, F. E.: A radiographic and histologic study on reattachment. J Periodontol 23: 158, 1952.
578. Arrocha, R., Wittwer, J. W., and Gargiulo, A. W.: Tissue response to heterogenous bone implantation in dogs. J Periodontol 39: 162, 1968.
579. Emmings, F. G.: Chemically modified osseous material for the restoration of bone defects. J Periodontol 45: 385, 1974.
580. Boyne, P. J., and Lyon, H. W.: Long-term histologic response to implants of anorganic heterogenous bone in man. J Dent Res 38: 699, 1959. IADR Abstr No. S15.

581. Melcher, A. H.: The use of heterogenous anorganic bone as an implant material in oral procedures. Oral Surg 15: 996, 1962.
582. Nielsen, I. M., Ellegaard, B., and Karring, T.: Kiel bone in healing interradicular lesions in monkeys. J Periodont Res 15: 328, 1980.
583. Hiatt, W. H., Solomons, C. C., and Butler, E. D.: The induction of new bone and cementum formation. II. Utilizing a collagen extract of ox bone. J Periodontol 41: 274, 1970.
584. Skinner, R. A., Toto, P. D., and Gargiulo, A. W.: Xenogeneic implants in primates. Collagen and chondroitin sulfate. J Periodontol 47: 196, 1976.
585. Solomons, C., and Gregory, G. W.: An evaluation of the effects of collagen implants on new bone formation in vivo. J Periodont Res 1: 218, 1966.
586. Shamash, D. B., and Bissada, N. F.: Stimulation of osteogenesis by topical application of calcitonin in periodontal osseous defects. J Periodontol 48: 619, 1977.
587. Packer, M. P., and Schaffer, E. M.: Cementum-dentine and Gelfoam implants in surgical periodontal pockets in dogs. Oral Surg 18: 722, 1964.
588. Schaffer, E. M.: Cementum and dentin implants in a dog and a rhesus monkey. J Periodontol 28: 125, 1957.
589. Stallard, R. E., and Hiatt, W. H.: The induction of new bone and cementum formation. I. Retention of mineralized fragments within the flap. J Periodontol 39: 273, 1968.
590. Schaffer, E. M.: Cartilage grafts in human periodontal pockets. J Periodontol 29: 176, 1958.
591. Held, A-J., and Spirgi, M.: Treatment of complex parodontitis by insertion of lyophilized cartilaginous grafts. Parodontologie 14: 9, 1960.
592. Schaffer, E. M.: Cartilage transplants into the periodontium of rhesus monkeys. Oral Surg 9: 1233, 1956.
593. Klingsberg, J.: Preserved sclera in periodontal surgery. J Periodontol 43: 634, 1972.
594. Klingsberg, J.: Periodontal scleral grafts and combined grafts of sclera and bone: Two-year appraisal. J Periodontol 45: 262, 1974.
595. Feingold, J. P., Chasens, A. I., Doyle, J., and Alfano, M. C.: Preserved scleral allografts in periodontal defects in man. II. Histologic evaluation. J Periodontol 48: 4, 1977.

596. Iacono, V. J., Gomes, B. C., Sciubba, J. J., Davis, R. K., Klingsberg, J., and Baer, P. N.: Clinical and laboratory studies on human sclera allografts. J Periodontol 51: 211, 1980.
597. Radentz, W. H., and Collings, C. K.: The implantation of plaster of Paris in the alveolar process of the dog. J Periodontol 36: 357, 1965.
598. Shaffer, C. D., and App, G. R.: The use of plaster of Paris in treating infrabony periodontal defects in humans. J Periodontol 42: 685, 1971.
599. Alderman, N. E.: Sterile plaster of Paris as an implant in the infrabony environment: A preliminary study. J Periodontol 40: 11, 1969.
600. Levin, M. P., Getter, L., Cutright, D. E., and Bhaskar, S. N.: Biodegradable ceramic in periodontal defects. Oral Surg 38: 344, 1974.
601. Levin, M. R., Getter, L., and Cutright, D. E.: A comparison of iliac marrow and biodegradable ceramic in periodontal defects. J Biomed Mater Res 9: 183, 1975.
602. Nery, E. B., Lynch, K. L., Hirthe, W. M., and Mueller, K. H.: Bioceramic implants in surgically produced infrabony defects. J Periodontol 46: 328, 1975.
603. Nery, E. B., and Lynch, K. L.: Preliminary clinical studies of bioceramic in periodontal osseous defects. J Periodontol 49: 523, 1978.
604. Bump, R. L., Salimeno, T., Hooker, S. P., Wilinison, E. G.: The use of woven ceramic fabric as a periodontal allograft. J Periodontol 46: 453, 1975.
605. Radell, R. L., and Cassingham, R. J.: A clinical evaluation of proplast as a periodontal implant material. J Periodontol 51: 110, 1980.
606. Ellegaard, B., Karring, T., Listgarten, M. and L  e, H.: New attachment after treatment of interradicular lesions. J Periodontol 44: 209, 1973.
607. Ellegaard, B., Karring, T., and L  e, H.: The fate of vital and devitalized bone grafts in the healing of interradicular lesions. J Periodont Res 10: 88, 1975.
608. Nilv  us, R., Johansson, O., and Egelberg, J.: The effect of autogenous cancellous bone grafts on healing of experimental furcation defects in dogs. J Periodont Res 13: 532, 1978.
609. Nilv  us, R., Bogle, G., Crigger, M., Egelberg, J., and Selvig, K. A.: The effect of topical citric acid application on the healing of experimental furcation defects in dogs. II. Healing after repeated surgery. J Periodont Res 15: 544, 1980.

610. Nilvéus, R., and Egelberg, J.: The effect of topical citric acid application on the healing of experimental furcation defects in dogs. III. The relative importance of coagulum support, flap design and systemic antibiotics. J Periodont Res 15: 551, 1980.
611. Ellegaard, B.: Bone grafts in periodontal attachment procedures. J Clin Periodontol 3: Extra Issue, 1976.
612. Schallhorn, R. G.: Present status of osseous grafting procedures. J Periodontol 48: 570, 1977.
613. Shellow, R. A., and Ratcliff, P. A.: The problems of attaining new alveolar bone after periodontal surgery. Periodont Abstr 15: 154, 1967.
614. Wade, A. B.: An evaluation of reparative procedures in periodontal surgery. Internat Dent J 21: 45, 1971.
615. Zander, H. A., Polson, A. M., and Heil, L.: Goals of periodontal therapy. J Periodontol 47: 261, 1976.
616. McDonald, F. L., Saari, T. T., and Davis, R. L.: A comparison of tooth accumulated materials from quadrants treated surgically or nonsurgically. J Periodontol 48: 147, 1977.
617. Ramfjord, S. P., Knowles, J. W., Nissle, R. R., Burgett, E. G., and Shick, R. A.: Results following three modalities of periodontal therapy. J Periodontol 46: 522, 1975.
618. Ramfjord, S. P., and Ash, M. M. Jr.: Periodontology and Periodontics. p. 608. Philadelphia, W. B. Saunders, 1979.
619. Leonard, H. J.: Raising the line of epithelial attachment and increasing depth of clinical root. J Periodontol 21: 221, 1950.
620. Weeks, P. R.: Pros and cons of periodontal pocket elimination procedures. Periodont Abstr 28: 4, 1980.
621. Alexander, A. G.: The effect of subgingival scaling on gingival inflammation. J Periodontol 40: 717, 1969.
622. Løe, H., Theilade, E., and Borglum-Jensen, S.: Experimental gingivitis in man. J Periodontol 36: 177, 1965.
623. Theilade, E., Wright, W. H., Borglum-Jensen, S., and Løe, H.: Experimental gingivitis in man. II. A longitudinal clinical and bacteriological investigation. J Periodont Res 1: 1, 1966.
624. Ringle, L. L.: A comparison of pocket reduction following hand and ultrasonic instrumentation. Periodont Abstr 15: 57, 1967.

625. Suomi, J. D., Green, J. C., Vermillion, J. R., Doyle, J., Chang, J., and Leatherwood, E. C.: The effect of controlled oral hygiene procedures on the progression of periodontal disease in adults: Results after third and final year. J Periodontol 42: 152, 1971.
626. Torafson, T., Kiger, R., Selvig, K. A., and Egelberg, J.: Clinical improvement of gingival conditions following ultrasonic versus hand instrumentation of periodontal pockets. J Clin Periodontol 6: 165, 1979.
627. Morrison, E. C., Lang, N. P., L  e, H., and Ramfjord, S. P.: Effects of repeated scaling and root planing and/or controlled oral hygiene on the periodontal attachment level and pocket depth in beagle dogs. I. Clinical findings. J Periodont Res 14: 428, 1979.
628. Badersten, A., Nilv  us, R., and Egelberg, J.: Effect of non-surgical periodontal therapy. I. Moderately advanced periodontitis. J Clin Periodontol 7: 336, 1980. Abstract.
629. Tagge, D. L., O'Leary, T. J., and El-Kafrawy, A. H.: The clinical and histological responses of periodontal pockets to root planing and oral hygiene. J Periodontol 46: 527, 1975.
630. Hughes, T. P., and Caffesse, R. G.: Gingival changes following scaling, root planing and oral hygiene. J Periodontol 49: 245, 1978.
631. Morrison, E. C., Ramfjord, S. P., and Hill, R. W.: Short-term effects of initial nonsurgical periodontal treatment (hygienic phase). J Clin Periodontol 7: 199, 1980.
632. Waite, I. M.: A comparison between conventional gingivectomy and a non-surgical regime in the treatment of periodontitis. J Clin Periodontol 3: 173, 1976.
633. Ramfjord, S.: Scaling and curettage -- is it enough? Periodont Abstr 27: 126, 1979.
634. Lang, N. P., Morrison, E. C., L  e, H., and Ramfjord, S. P.: Longitudinal therapeutic effects on the periodontal attachment level and pocket depth in beagle dogs. J Periodont Res 14: 418, 1979.
635. Pihlstrom, B. L., and Ortiz, C.: Four year longitudinal study of two methods of periodontal therapy. J Dent Res 59 (Special Issue A): 298, 1980. AADR Abstr No. 122.
636. Knowles, J. W.: Oral hygiene related to long-term effects of periodontal therapy. Mich Dent A J 55: 147, 1973.

637. Nyman, S., Rosling, B., and Lindhe, J.: Effect of professional tooth cleaning on healing after periodontal surgery. J Clin Periodontol 2: 80, 1975.
638. Nyman, S., Lindhe, J., and Rosling, B.: Periodontal surgery in plaque-infected dentitions. J Clin Periodontol 4: 240, 1977.
639. Polson, A. M., Kantor, M. E., and Zander, H. A.: Periodontal repair after reduction of inflammation. J Periodont Res 14: 520, 1979.
640. Kantor, M.: The behavior of angular bony defects following reduction of inflammation. J Periodontol 51: 433, 1980.
641. Bogle, G., Rathbun, E., Oliver, R., Hornbuckle, C., and Egelberg, J.: Effect of post-operative use of chlorhexidine on regeneration of bifurcation defects in dogs. J Periodont Res 9: 127, 1974.
642. Hugoson, A., and Schmidt, G.: Influence of plaque control on the healing of experimentally-induced bone defects in the dog. J Periodontol 49: 135, 1978.
643. Bell, D. C., and Rule, R. W. Jr.: The periodontal lesion. II. A roentgenographic evaluation of treated periodontal cases. J Periodontol 21: 70, 1950.
644. Lovdal, A., Arno, A., Schei, O., and Waerhaug, J.: Combined effect of subgingival scaling and controlled oral hygiene on the incidence of gingivitis. Acta Odont Scand 19: 537, 1961.
645. Oliver, R. C.: Tooth loss with and without periodontal therapy. Periodont Abstr 17: 8, 1969.
646. Ross, I. F., and Thompson, R. H.: The results of treatment. A long term study of one hundred and eighty patients. Parodontologie 25: 125, 1971.
647. Lindhe, J., and Nyman, S.: The effect of plaque control and surgical pocket elimination on the establishment and maintenance of periodontal health. A longitudinal study of periodontal therapy in cases of advanced disease. J Clin Periodontol 2: 67, 1975.
648. Hirschfeld, L., and Wasserman, B.: A long-term survey of tooth loss in 600 treated periodontal patients. J Periodontol 49: 225, 1978.
649. Ross, I. F., and Thompson, R. H.: A long term study of root retention in the treatment of maxillary molars with furcation involvement. J Periodontol 49: 238, 1978.
650. Feinstein, A. R.: Clinical Judgement. Baltimore, Williams and Wilkins, 1967.




Fig. 1. Dotted line represents cut of internal gingival incision leaving relatively constant connective tissue gingival wound surface between epithelium and alveolar process.




Fig. 2. After removal of a wedge of tissue note that tissue remnants remain in areas of the epithelial attachment and the connective tissues of zones 3 and 4.



Figure 3. Histologic section of tooth and gingiva, immediately after internal bevel incision and removal of pocket lining to the depth of probing. (Rhesus monkey, H and E stain, original magnification X10).

ACKNOWLEDGEMENTS

The author is indebted to Dr. Perry Ratcliff and to Dr. Fermin Carranza, Jr., Editor of the Journal of the Western Society of Periodontology, for permission to use Fig. 1 and Fig. 2 from Periodont Abstr 14: 57, 1966. The exceptionally fine work of manuscript preparation was the contribution of Ms. Myra Rouse and Syd Kline.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO. AD-A095 907	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A COMPREHENSIVE REVIEW OF NEW ATTACHMENT THERAPY.		5. TYPE OF REPORT & PERIOD COVERED
7. AUTHOR(s) M. R. WIRTHLIN, Jr.		6. PERFORMING ORG. REPORT NUMBER NDRI-PR-81-01
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Dental Research Institute Naval Base, Bldg. 1-H Great Lakes, IL 60088		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Medical Research and Development Command National Naval Medical Center Bethesda, MD 20014		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 63706N M0095PN003.3010
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Bureau of Medicine and Surgery Department of the Navy Washington, D. C. 20372		12. REPORT DATE JANUARY 1981
		13. NUMBER OF PAGES 76
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) This document has been approved for public release; distribution unlimited.		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Dentistry Regeneration (physiology) Surgery Healing Epithelium Connective tissue		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Dental surgical procedures have developed over the years based on strong clinical impression and empirical practices. Important advances have been made in the field of clinical trials both in measurement of conditions and in analysis of clinical data. Many articles in the past have dwelled upon methods of handling soft tissues and bony defects, while recently there has been interest in the treatment of the diseased root surface. This report is a review of the literature in 1980, and highlights the biological nature of the healing dentogingival junction.		


DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S. N 0102- LF-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)



S/N 0102- LF-014-6601

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)